

A new genus and species of Platyischnopidae (Amphipoda: Gammaridea) from the Argentine Sea, South-West Atlantic Ocean

IGNACIO L. CHIESA^{1,2,3} & GLORIA M. ALONSO¹

¹Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, División Invertebrados, Av. Ángel Gallardo 470, C1405DJR, Buenos Aires, Argentina

²Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Biodiversidad y Biología Experimental, C1428HA, Buenos Aires, Argentina

³Corresponding author. E-mail: ichiesa@macn.gov.ar

Abstract

The family Platyischnopidae is herein reported for the first time in the Argentine Sea, South-West Atlantic Ocean. A new genus and species, *Platyisao holodividum gen. et. sp. nov.*, collected off the coast of Buenos Aires and Río Negro provinces, is fully described and illustrated. *Platyisao* gen. nov. is distinguished from the eight other genera of Platyischnopidae by the gnathopods subchelate, and the telson elongate, completely cleft. In addition, the distribution of *Tiburonella viscana* (Barnard J.L., 1964), up to now known in the South-West Atlantic Ocean from Brazilian waters, is extended to the coast off Buenos Aires province, Argentina.

Key words: Amphipods, platyischnopids, Southwest Atlantic, Argentina, *Platyisao holodividum* gen. et. sp. nov., *Tiburonella viscana* distribution

Introduction

Platyischnopidae Barnard & Drummond, 1979 is a gammaridean benthic family with characteristics typical of fossorial amphipods, such as an elongate cephalothorax and antennae, pereopods and uropods with many setae and spines (Thomas & Barnard 1983). Platyischnopids are fast swimmers and efficient predators, abundant on soft bottoms of the continental shelves, and are also frequent in reef clasts and on seagrass beds, at depths ranging from intertidal to 175 m; most species inhabit warm and temperate waters (Thomas 1993; Hughes 2009; Souza-Filho & Serejo 2012; and Table 1).

The genus *Platyischnopus* Stebbing, 1888 was originally included in Pontoporeiidae by Stebbing (1888) and then alternatively assigned to Phoxocephalidae or Haustoriidae (Chilton 1922; Oliveira 1955; Bousfield 1970; Rabindranath 1971; Griffiths 1974a,b; Ledoyer 1979, among others). The family Platyischnopidae was erected by Barnard & Drummond (1979) with *Platyischnopus* as the type genus. Platyischnopids were distinguished from haustoriids and phoxocephalids principally by the shape of the rostrum, the mandibular molar and the pereopod 7 (Barnard & Drummond 1979), and can be easily distinguished from other amphipod families by the cone-shaped rostrum covered with apical sensory pits (Hughes 2009).

At present, the genus *Platyischnopus* includes two species: *P. mam* Barnard & Drummond, 1979 and *P. mirabilis* Stebbing, 1888, with both species distributed in Australian waters (Barnard & Drummond 1979; Hughes 2009).

Besides *Platyischnopus*, other four genera were also erected in Platyischnopidae by Barnard & Drummond (1979): *Indischnopus*, *Tittakunara*, *Tomituka*, and *Yurrokus*.

The genus *Indischnopus* was established to accommodate *I. herdmani* (Walker, 1904) distributed in the Indic Ocean and *I. capensis* (Barnard K.H., 1925) from South Africa, the two species being originally placed in *Platyischnopus* (Barnard & Drummond 1979). Othman & Morino (1996) reported *I. redangi* from Malaysian waters.

anonymous reviewer for providing constructive comments. This research was partially supported by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET PIP 11220090100375) and the Universidad de Buenos Aires (UBACyT 20020100100857).

References

- Abessa, D.M.S. & Sousa, E.C.P.M. (2003) Sensitivity of the amphipod *Tiburonella viscana* (Platyischnopidae) to K₂Cr₂O₇. *Brazilian Archives of Biology and Technology*, 46, 53–55.
<http://dx.doi.org/10.1590/s1516-89132003000100009>
- Abessa, D.M.S., Carr, R.S., Rachid, B.R.F., Sousa, E.C.P.M., Hortelani, M.A. & Sarkis, J.E. (2005) Influence of a Brazilian sewage outfall on the toxicity and contamination of adjacent sediments. *Marine Pollution Bulletin*, 50, 875–885.
<http://dx.doi.org/10.1016/j.marpolbul.2005.02.034>
- Abessa, D.M.S., Sousa, E.C.P.M., Rachid, B.R.F. & Mastroti, R.R. (1998) Use of the burrowing amphipod *Tiburonella viscana* as a tool in marine sediments contamination assessment. *Brazilian Archives of Biology and Technology*, 41 (2), 225–230.
- Atienza, A. & Martín, A. (2001) *Tiburonella viscana* (Amphipoda: Platyisnopidae) en las costas de Venezuela. *Revista de Biología Tropical*, 49 (3), 1270.
- Balech, E. & Ehrlich, M.D. (2008) Esquema biogeográfico del Mar Argentino. *Revista de Investigación y Desarrollo Pesquero*, 19, 45–75.
- Barnard, J.L. (1964) Los anfípodos bentónicos marinos de la costa occidental de Baja California. *Revista de la Sociedad Mexicana de Historia Natural*, 24, 205–274.
- Barnard, J.L. & Drummond, M.M. (1979) Gammaridean Amphipoda of Australia, Part IV. *Smithsonian Contributions to Zoology*, 269, 1–69.
- Barnard, J.L. & Karaman, G.S. (1991) The families and genera of marine gammaridean Amphipoda (except marine gammaroids). *Records of the Australian Museum*, supplement 13 (Part 2), 419–866.
<http://dx.doi.org/10.3853/j.0812-7387.13.1991.367>
- Barnard, K.H. (1925) Contributions to the Crustacean Fauna of South Africa, Number 8: Further Additions to the List of Amphipoda. *Annals of the South African Museum*, 20, 319–330.
- Bousfield, E.L. (1970) Adaptive Radiation in Sand-burrowing Amphipod Crustaceans. *Chesapeake Science*, 11 (3), 143–154.
<http://dx.doi.org/10.2307/1351237>
- Cesar, A., Abessa, D.M.S., Pereira, C.D.S., Santos, A.R., Fernández, N., Choueri, R.B. & Delvalls, T.A. (2009) A simple approach to integrate the ecotoxicological and chemical data for the establishment of environmental risk. *Brazilian Archives of Biology and Technology*, 52 (1), 233–240.
<http://dx.doi.org/10.1590/s1516-89132009000100028>
- Chilton, C. (1922) Amphipoda: results of Dr. E. Mjobergs Swedish Scientific Expeditions to Australia 1910–13, XXXI. *Kungliga Svenska Vetenskapsakademiens Handlingar*, 63 (3), 1–11.
- Coleman, C.O. (2003) Digital inking: How to make perfect line drawings on computers. *Organisms Diversity & Evolution*, 3, (Electr. Suppl. 14), 1–14. Available from: <http://senckenberg.de/odes/03-14.htm> (accessed 13 Feb 2013)
- Da Silva, P.S.R., Das Neves, L.P. & Bemvenuti, C.E. (2008) Temporal variation of sandy beach macrofauna at two sites with distinct environmental conditions on Cassino beach, extreme southern Brazil. *Brazilian Journal of Oceanography*, 56 (4), 257–270.
<http://dx.doi.org/10.1590/s1679-87592008000400001>
- Das Neves, L.P., Da Silva, P.S.R. & Bemvenuti, C.E. (2008) Temporal variability of benthic macrofauna on Cassino beach, southernmost Brazil. *Iheringia, Série Zoologia*, 98 (1), 36–44.
<http://dx.doi.org/10.1590/s0073-47212008000100005>
- Granda, L.D., Fockedey, N., De Mey, M., Beyst, B., Cornejo, M.D.P., Calderon, J. & Vincx, M. (2004) Spatial patterns of the surf zone hyperbenthic fauna of Valdivia Bay (Ecuador). *Hydrobiologia*, 529, 205–224.
<http://dx.doi.org/10.1007/s10750-004-6417-3>
- Griffiths, C.L. (1974a) The Amphipoda of Southern Africa. Part 3. The Gammaridea and Caprellidea of Natal. *Annals of the South African Museum*, 62 (7), 209–264.
- Griffiths, C.L. (1974b) The Amphipoda of Southern Africa. Part 4. The Gammaridea and Caprellidea of the Cape Province East of Cape Agulhas. *Annals of the South African Museum*, 65 (9), 251–336.
- Guerrero, R.A. & Piola, A.R. (1997) Masas de agua en la Plataforma Continental. In: Boschi, E.E. (Ed.), *El Mar Argentino y sus recursos pesqueros. Tomo I. Antecedentes históricos de las exploraciones en el mar y las características ambientales*. Publicaciones especiales INIDEP, Mar del Plata, Argentina, pp. 107–118.
- Hughes, L.E. (2009) Platyischnopidae. In: Lowry, J.K. & Myers, A.A. (Eds.), Benthic Amphipoda of the Great Barrier Reef, Australia. *Zootaxa*, 2260, 828–835.
- Ledoyer, M. (1979) Les gammariens de la pente externe du grand récif de Tuléar (Madagascar) (Crustacea Amphipoda). *Memorie del Museo Civico di Storia Naturale di Verona*, Series 2, Sezione Scienze della Vita, 2, 1–150.
- López Gappa, J., Alonso, G.M. & Landoni, N.A. (2006) Biodiversity of benthic Amphipoda (Crustacea: Peracarida) in the

- Southwest Atlantic between 35°S and 56°S. *Zootaxa*, 1342, 1–66.
- Maranho, L.A., Abreu, I., Santelli, R., Cordeiro, R.C., Soares-Gomes, A., Moreira, L.B., Morais, R.D. & Abessa, D.M.S. (2009) Sediment toxicity assessment of Guanabara Bay, Rio de Janeiro, Brazil. *Journal of Coastal Research* (Special Issue), 56, 851–855.
- Martinez, M.I. & Brogger, M.I. (2012) *Thandarum hernandezi*, a new genus and new species of sea cucumber family Sclerodactylidae (Echinodermata: Holothuroidea: Dendrochirotida) from the Southwestern Atlantic Ocean. *Zootaxa*, 3304, 63–68
- Melo, S.L.R. & Nipper, M. (2007) Sediment toxicity tests using the burrowing amphipod *Tiburonella viscosa* (Amphipoda: Platyischnopidae). *Ecotoxicology and Environmental Safety*, 66, 412–420.
<http://dx.doi.org/10.1016/j.ecoenv.2005.12.003>
- Oliveira, L.P.H. (1955) *Phoxocephalus capuciatus*, nova espécie de Crustacea Amphipoda, Phoxocephalidae. *Memorias do Instituto Oswaldo Cruz*, 53 (2–4), 313–319.
<http://dx.doi.org/10.1590/s0074-02761955000200009>
- Ortiz, M., Martín, A. & Atienza, D. (2000) Una nueva especie de anfípodo del género *Tiburonella* (Crustacea, Amphipoda, Platyischnopidae) de Venezuela. *Acta Biológica Venezolica*, 20 (2), 29–36.
- Ortiz, M., Martín, A. & Díaz, Y.J. (2007) Lista y referencias de los crustáceos anfípodos (Amphipoda: Gammaridea) del Atlántico occidental tropical. *Revista de Biología Tropical*, 55 (2), 479–528.
- Othman, B.H.R. & Morino, H. (1996) A new species of the genus *Indischmopus* from Malaysia (Crustacea, Amphipoda, Platyischnopidae). *Bollettino del Museo Civico di Storia Naturale di Verona*, 20, 105–115.
- Piola, A.R. & Rivas, A.L. (1997) Corrientes en la Plataforma Continental. In: Boschi, E.E. (Ed.), *El Mar Argentino y sus recursos pesqueros. Tomo I. Antecedentes históricos de las exploraciones en el mar y las características ambientales*. Publicaciones especiales INIDEP, Mar del Plata, Argentina, pp. 119–132.
- Rabindranath, P. (1971) Haustoriid Amphipods (Crustacea) from India. *Hydrobiologia*, 38, 521–839.
<http://dx.doi.org/10.1007/bf00036557>
- Roccatagliata, D., Chiesa, I.L., Raffo, M.P. & Gómez Simes, E. (2010) On the occurrence of the genus *Paranebalia* (Crustacea: Phyllocarida:Leptostraca) in northern Patagonia, Argentina. *Zootaxa*, 2349, 65–68.
- Ríos, C., Mutschke, E. & Morrison, E. (2003) Biodiversidad bentónica sublitoral en el estrecho de Magallanes, Chile. *Revista de Biología Marina y Oceanografía*, 38 (1), 1–12.
- Schellenberg, A. (1931) Gammariden und Caprelliden des Magellangebietes, Südgeorgiens und der Westantarktis. *Further Zoological Results of the Swedish Antarctic Expedition 1901–1903*, 2 (6), 1–290.
- Shoemaker, C.R. (1942) Amphipod crustaceans collected on the Presidential Cruise of 1938. *Smithsonian Miscellaneous Collections*, 101 (11), 1–52.
- Souza-Filho, J.F. & Serejo, C.S. (2012) Redescription and designation of a neotype for *Eudevenopus capuciatus* (Oliveira, 1955) (Crustacea: Amphipoda: Platyischnopidae) from Brazilian waters, with comments on its cuticular ultrastructures. *Cahiers de Biologie Marine*, 43, 469–484.
- Stebbing, T.R.R. (1888) Report on the Amphipoda collected by H. M. S. Challenger during the years 1873–1876. *Report on the Scientific Results of the Voyage of H.M.S. Challenger during the years 1873–76*, 29 (67), 1–1737.
- Thomas, J.D. (1993) *Identification manual for marine Amphipoda: (Gammaridea) I. Common coral reef and rocky bottom amphipods of South Florida*. State of Florida, Department of Environmental Protection, Tallahassee, Florida, USA, 83 pp.
- Thomas, J.D. & Barnard, J.L. (1983) The Platyischnopidae of America (Crustacea: Amphipoda). *Smithsonian Contributions to Zoology*, 375, 1–33.
- Torres, R.J., Abessa, D.M.S., Santos, F.C., Maranho, L.A., Davanco, M.B., Do Nascimento, M.R.L. & Mozeto, A.A. (2009) Effects of dredging operations on sediment quality: contaminant mobilization in dredged sediments from the Port of Santos, SP, Brazil. *Journal of Soils and Sediments*, 9, 420–432.
<http://dx.doi.org/10.1007/s11368-009-0121-x>
- Wakabara, Y. (1969) Sobre alguns Gammaridea (Crustacea-Amphipoda) da região de Ubatuba. Dissertação de mestrado. Universidade de São Paulo, Instituto Oceanográfico, São Paulo, Brazil, 52 pp.
- Wakabara, Y., Taraman, A.S., Valério-Berardo, M.T., Duleba, W. & Pereira Leite, F.P. (1991) Gammaridean and caprellidean fauna from Brazil. *Hydrobiologia*, 223, 69–77.
<http://dx.doi.org/10.1007/bf00047629>
- Wakabara, Y. & Serejo, C.S. (1998) Malacostraca–Peracarida. Amphipoda Gammaridea and Caprellidea. In: Young, P.S. (Ed.), *Catalogue of Crustacea of Brazil*, Série Livros n. 6. Museu Nacional, Rio de Janeiro, Brazil, pp. 561–594.
- Walker, A.O. (1904) Report on the Amphipoda Collected by Professor Herdman, at Ceylon, in 1902. Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar, Supplementary reports, 17, 229–300.
- Watling, L. (1989) A classification system for crustacean setae based on the homology concept. In: Felgenhauer, B.E., Watling, L. & Thistle, A.B. (Eds.), *Functional morphology of feeding and grooming in Crustacea*. Crustacean issues 6, Balkema, A.A., Rotterdam, The Netherlands, pp. 15–26.
- Young, P.S. & Serejo, C.S. (2005) Crustacea of the Abrolhos Region, Brazil. In: Dutra, G.F., Allen, G.R., Werner, T. & McKenna, S.A. (Eds.), *A rapid marine biodiversity assessment of the Abrolhos Bank, Bahia, Brazil*. RAP Bulletin of Biological Assessment 38, Conservation International, Washington, USA, pp. 91–95.