

A new species of *Pseudopaludicola* Miranda-Ribeiro, 1926 (Anura: Leptodactylidae: Leiuperinae) from northwestern state of São Paulo, Brazil

ANDRÉ PASONATO^{1,7}, JESSICA RHAIZA MUDREK², ANA CRISTINA PRADO VEIGA-MENONCELLO³, DENISE DE CERQUEIRA ROSSA-FERES⁴, ITAMAR ALVES MARTINS^{1,5} & CHRISTINE STRÜSSMANN^{2,6}

¹Pós-Graduação em Biologia Animal, Universidade Estadual Paulista (UNESP), Rua Cristóvão Colombo, 2265, Jardim Nazareth, 15054–000, São José do Rio Preto, São Paulo, Brazil

²Pós-Graduação em Ecologia e Conservação da Biodiversidade, Instituto de Biociências, Universidade Federal de Mato Grosso (UFMT), Instituto de Biociências, Universidade Federal de Mato Grosso (UFMT), Av. Fernando Correia da Costa, 2367, Boa Esperança, 78060–900, Cuiabá, Mato Grosso, Brazil

³Departamento de Biologia Estrutural e Funcional, Instituto de Biologia, Universidade Estadual de Campinas (UNICAMP), 13083–863, Campinas, São Paulo, Brazil

⁴Departamento de Zoologia e Botânica, Universidade Estadual Paulista (UNESP), Rua Cristóvão Colombo, 2265, Jardim Nazareth, 15054–000, São José do Rio Preto, São Paulo, Brazil

⁵Laboratório de Zoologia, Instituto Básico de Biociências, Universidade de Taubaté (UNITAU), Av. Tiradentes, 500, 12030-180, Taubaté, São Paulo, Brazil

⁶Departamento de Ciências Básicas e Produção Animal, Faculdade de Agronomia, Medicina Veterinária e Zootecnia, Universidade Federal de Mato Grosso (UFMT), Av. Fernando Correia da Costa, 2367, Boa Esperança, 78060–900, Cuiabá, Mato Grosso, Brazil

⁷Corresponding author. E-mail: andreufmt@gmail.com

Abstract

A new species of *Pseudopaludicola* is described from human-altered areas originally covered by Semideciduous Forest in northwestern state of São Paulo, southeastern Brazil. Morphologically, the new species differs from four species belonging to the *P. pusilla* group by the absence of either T-shaped terminal phalanges or toe tips expanded, and from all other congeners except *P. canga* and *P. facureae* by possessing an areolate vocal sac, with dark reticulation. The higher duration (300–700 ms) of each single, pulsed note (9–36 nonconcatenated pulses) that compose the call in the new species distinguishes it from all other 14 species of *Pseudopaludicola* with calls already described (10–290 ms). Absence of harmonics also differ the advertisement call of the new species from the call of its sister species *P. facureae*, even though these two species presented unexpected low genetic distances. Although we could not identify any single morphological character distinguishing the new species from *P. facureae*, a PCA and DFA performed using 12 morphometric variables evidenced significant size differences between these two species.

Key words: Amphibia, taxonomy, advertisement call, genetic distance

Introduction

Species from the genus *Pseudopaludicola* Miranda-Ribeiro, 1926 are distributed in open grasslands (seasonally flooded or not), open environments in the vicinities or amidst tropical rainforests, and dry forests that occur throughout South America (Lynch 1989; Pansonato *et al.* 2012a; Frost 2014). The genus currently comprises 17 valid species (Frost 2014; Pansonato *et al.* 2014).

According to former phylogenetic analyses by Lobo (1995) and by Cardozo and Suárez (2012), only the *Pseudopaludicola pusilla* group, characterized by T-shaped terminal phalanges was recognized as monophyletic. Four species of *Pseudopaludicola* were assigned to this group: *P. boliviana* Parker, 1927; *P. ceratophyes* Rivero & Serna, 1984; *P. llanera* Lynch, 1989; and *P. pusilla* (Ruthven, 1916). Until recently, other 14 species remained unassigned to this or to any other monophyletic group (Lynch 1989; Cardozo & Suárez 2012; Andrade & Carvalho, 2013; Roberto *et al.* 2013; Magalhães *et al.* 2014); *P. ameghini* (Cope, 1887); *P. canga* Giaretta & Kokubum, 2003;

Geographical, morphological, ecological, bioacoustic, and genetic patterns of co-occurring species of *Pseudopaludicola* should be evaluated using integrative approaches, for a better understanding of habitat use, geographic distribution, and evolution in the genus.

TABLE 5. Measurements of acoustic parameters of specimens of *Pseudopaludicola facureae* from the type locality (Uberlândia, Minas, Brazil). Values are: mean \pm standard deviation (minimum–maximum). N: number of calls (= notes) analyzed (number of specimens recorded).

Advertisement call parameters	<i>Pseudopaludicola facureae</i> N=280 (8)
Number of notes per series (s)	15.8 \pm 10.1 (2–55)
Series duration (s)	0.8 \pm 0.6 (0.1–4.3)
Inter-series interval (s)	0.5 \pm 0.2 (0.2–1.7)
Note duration (s)	0.02 \pm 0.005 (0.01–0.04)
Inter-notes interval (s)	0.03 \pm 0.03 (0.01–0.4)
Minimum frequency (Hz)	2688.5 \pm 573.9 (1114.3–3992.6)
Maximum frequency (Hz)	5947.9 \pm 336.8 (5372.1–6771.4)
Dominant frequency (Hz)	4559.4 \pm 212.2 (4306.6–4995.7)
Harmonic 1 (Hz)	4306–4995
Harmonic 2 (Hz)	8806–9495
Harmonic 3 (Hz)	13306–13995
Average rate of notes/min	1249.1 \pm 388.5 (646–2000)

Acknowledgements

Special thanks go to the owner of Boa Esperança smallholding, Toninho, for allowing our field work there; to Miguel Vences, for comments on earlier versions of the manuscript; to Felipe Curcio, for allowing access to material under his care at “Coleção Zoológica de Vertebrados da Universidade Federal de Mato Grosso” (UFMT); to Ariovaldo Giaretta and Felipe de Andrade for allowing access to topotypic calls and measurements of *P. facureae*; to CNPq (project “Rede de pesquisa em anfíbios e répteis de ecossistemas não florestais brasileiros”; process # 563352/2010-8) and to Evandson Silva (FAPEMAT process # 285060/2010), for financial and logistical support, respectively. To Rodrigo Silva for help during field sampling; to Jéssica Cantarini for clearing and staining specimens; to Jeffrey Himmelstein for language review. Ecological information about habitat use was obtained during several research projects accomplished by students from the Theoretical Ecology Laboratory at UNESP to whom we are grateful: Rodolfo Pelinson, Rodrigo Silva, Michel Garey, Thiago Pereira, Veronica Sousa and Tales Cordeiro. AP thanks “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior” (CAPES) for a PhD scholarship; CS and DCRF thank to CNPq for research fellowships (processes # 309541/2012-3 and # 303522/2013-5).

References

- Andrade, F.S. & Carvalho, T.R. (2013) A new species of *Pseudopaludicola* Miranda-Ribeiro (Leiuperinae: Leptodactylidae: Anura) from the Cerrado of southeastern Brazil. *Zootaxa*, 3608, 389–397.
<http://dx.doi.org/10.11646/zootaxa.3608.5.7>
- Cardozo, D. & Suárez, P. (2012) Osteological description of *Pseudopaludicola canga* with implications for the taxonomic position of this taxon. *Zootaxa*, 3515, 75–82.
- Cardozo, D. & Toledo, L.F. (2013) Taxonomic status of *Pseudopaludicola riopiedadensis* Mercadal de Barrio and Barrio, 1994 (Anura, Leptodactylidae, Leiuperinae). *Zootaxa*, 3734 (5), 571–582.
<http://dx.doi.org/10.11646/zootaxa.3734.5.6>
- Carvalho, T.R. (2012) A new species of *Pseudopaludicola* Miranda-Ribeiro (Leiuperinae: Leptodactylidae: Anura) from the Cerrado of southeastern Brazil with a distinctive advertisement call pattern. *Zootaxa*, 3328, 47–54.
- De la Riva, I., Köhler, J., Lötters, S. & Reichle, S. (2000) Ten years of research on Bolivian amphibians: updated checklist,

- distribution, taxonomic problems, literature and iconography. *Revista Española Herpetologica*, 14, 19–164.
- Duarte, T.C., Veiga-Menoncello, A.C.P., Lima, J.F.R., Strüssmann, C., Del Grande, M.L., Giaretta, A.A., Pereira, E.G., Rossa-Feres, D.C. & Recco-Pimentel, S.M. (2010) Chromosome analysis in *Pseudopaludicola* (Anura, Leiuperidae), with description of sex chromosomes XX/XY in *P. saltica*. *Hereditas*, 147, 43–52.
<http://dx.doi.org/10.1111/j.1601-5223.2009.02153.x>
- Duellman, W.E. (1970) The Hylid Frogs of Middle America. *Monograph of the Museum of Natural History of University of Kansas*, 1, 1–753.
- Duré, M.I., Schaefer, E.F., Hamann, M.I. & Kehr, A.I. (2004) Consideraciones ecológicas sobre la dieta, la reproducción y el parasitismo de *Pseudopaludicola boliviensis* (Anura, Leptodactylidae) de Corrientes, Argentina. *Phyllomedusa*, 3, 121–131.
<http://dx.doi.org/10.11606/issn.2316-9079.v3i2p121-131>
- Fávero, E.R., Veiga-Menoncello, A.C.P., Rossa-Feres, D.C., Strüssmann, C., Giaretta, A.A., Andrade, G.V., Colombo, P. & Recco-Pimentel, S.M. (2011) Intrageneric karyotypic variation in *Pseudopaludicola* (Anura: Leiuperidae) and its taxonomic relatedness. *Zoological Studies*, 50, 826–836.
- Frost, D.R. (2014) *Amphibian Species of the World: an Online Reference*. Version 6.0 American Museum of Natural History, New York, USA. Available from: <http://research.amnh.org/herpetology/amphibia/> (accessed 16 April 2014)
- Giaretta, A.A. & Facure, K.G. (2009) Habitat, egg-laying behaviour, eggs and tadpoles of four sympatric species of *Pseudopaludicola* (Anura, Leiuperidae). *Journal of Natural History*, 43, 995–1009.
<http://dx.doi.org/10.1080/00222930802702456>
- Giaretta, A.A. & Kokubum, M.N.C. (2003) A new species of *Pseudopaludicola* (Anura, Leptodactylidae) from northern Brazil. *Zootaxa*, 383, 1–8.
- Haddad, C.F.B. & Cardoso, A.J. (1987) Taxonomia de três espécies de *Pseudopaludicola* (Anura, Leptodactylidae). *Papéis Avulsos de Zoologia*, 36, 287–300.
- Heyer, W.R., Rand, A.S., Cruz, C.A.G., Peixoto, O.L. & Nelson, C.E. (1990) Frogs of Boracéia. *Arquivos de Zoologia*, 31, 231–410.
- Jungfer, K.H., Faivovich, J., Padial, J.M., Castroviejo-Fisher, S., Lyra, M.M., Berneck, B.V.M., Iglesias, P.P., Kok, P.J.R., MacCulloch, R.D., Rodrigues, M.T., Verdade, V.K., Castello, C.P.T., Chaparro, J.C., Valdujo, P.H., Reichle, S., Moravec, J., Gvoždík, V., Gagliardi-Urritia, G., Ernst, R., De la Riva, I., Means, D.B., Lima, A.P., Señaris, C., Wheeler, W.C. & Haddad, C.F.B. (2013) Systematics of spiny-backed treefrogs (Hylidae: *Osteocephalus*): an Amazonian puzzle. *Zoologica Scripta*, 42, 351–380.
<http://dx.doi.org/10.1111/zsc.12015>
- Lavilla, E.O. & Cei, J.M. (2001) Amphibians of Argentina. A second update, 1987–2000. *Monografie Museo Regionale di Scienze Naturali Torino*, 28, 1–78.
- Lobo, F. (1994) Descripción de una nueva especie de *Pseudopaludicola* (Anura: Leptodactylidae), redescipción de *P. falcipes* (Hensel, 1867) y *P. saltica* (Cope, 1887) y osteología de las tres especies. *Cuadernos de Herpetología*, 8, 177–199.
- Lobo, F. (1995) Análisis filogenético del género *Pseudopaludicola* (Anura: Leptodactylidae). *Cuadernos de Herpetología*, 9, 21–43.
- Lobo, F. (1996) Evaluación del status taxonómico de *Pseudopaludicola ternetzi* Miranda Ribeiro, 1937; *P. mystacalis* y *P. ameghini* (Cope, 1887). Osteología y distribución de las especies estudiadas. *Acta Zoológica Lilloana*, 43, 327–346.
- Lötters, S., Schmitz, A., Reichle, S., Rödder, D., Quennet, V. (2009) Another case of cryptic diversity in poison frogs (Dendrobatidae: *Ameerega*) — description of a new species from Bolivia. *Zootaxa*, 2028, 20–30.
- Lynch, J.D. (1989) A review of the leptodactylid frogs of the genus *Pseudopaludicola* in northern South America. *Copeia*, 1989, 577–588.
<http://dx.doi.org/10.2307/1445483>
- Magalhães, F.M., Loebmann, D., Kokubum, M.N.C., Haddad, C.F.B. & Garda, A.A. (2014) A new species of *Pseudopaludicola* (Anura: Leptodactylidae: Leiuperinae) from northeastern Brazil. *Herpetologica*, 70, 77–88.
<http://dx.doi.org/10.1655/HERPETOLOGICA-D-13-00054>
- Martinelli, L.A. & Filoso, S. (2008) Expansion of sugar ethanol production in Brazil: environmental and social challenges. *Ecological Applications*, 18, 885–898.
<http://dx.doi.org/10.1890/07-1813.1>
- McLister, J.D., Stevens, E.D. & Bogart, J. (1995) Comparative contractile dynamics of calling and locomotor muscles in three hylid frogs. *Journal of Experimental Biology*, 198, 1527–1538.
- Miranda-Ribeiro, A. (1937) Alguns batrachios novos das coleções do Museu Nacional. *O Campo*, 66–69.
- Necchi Jr., O. (2011) *Fauna and flora of remnant forest fragments in the São Paulo state Northwest region. BIOTA-FAPESP Program*. FAPESP, São Paulo. Available from: http://media.fapesp.br/bv/uploads/publicacoes/pasta_biota_40.pdf (accessed 1 June 2013)
- Pansonato, A., Moraes, D.H., Ávila, R.W., Kawashita Ribeiro, R.A., Strüssmann, C. & Martins, I.A. (2012a) A new species of *Pseudopaludicola* Miranda-Ribeiro, 1926 (Anura: Leiuperidae) from the state of Mato Grosso, Brazil, with comments on the geographic distribution of *Pseudopaludicola canga* Giaretta & Kokubum, 2003. *Zootaxa*, 3523, 49–58.
- Pansonato, A., Mudrek, J.R., Rodrigues, T.F.D., Strüssmann, C. & Martins, I.A. (2012b) Ocorrência simpátrica de espécies do gênero *Pseudopaludicola* Miranda-Ribeiro, 1926 (Anura, Leiuperidae) em diferentes tipos de ambientes úmidos. *Anais do I Congresso Brasileiro de Áreas Úmidas - Água, Alimento, Energia, no Presente e no Futuro (I CONBRAU)*. CPP, Cuiabá, Mato Grosso, pp. 220–222.
- Pansonato, A., Strüssmann, C., Mudrek, J.R. & Martins, I.A. (2013) Morphometric and bioacoustic data on three species of

- Pseudopaludicola* Miranda-Ribeiro, 1926 (Anura: Leptodactylidae: Leiuperinae) described from Chapada dos Guimarães, Mato Grosso, Brazil, with the revalidation of *Pseudopaludicola ameghini* (Cope, 1887). *Zootaxa*, 3620 (1), 147–162.
<http://dx.doi.org/10.11646/zootaxa.3620.1.7>
- Pansonato, A., Mudrek, J.R., Simioni, F., Martins, I.A. & Strüssmann, C. (2014) Geographical variation in morphological and bioacoustic traits of *Pseudopaludicola mystacalis* (Cope, 1887) and a reassessment of the taxonomic status of *Pseudopaludicola serrana* Toledo, 2010 (Anura: Leptodactylidae: Leiuperinae). *Advances in Zoology*, 563165, 1–13.
<http://dx.doi.org/10.1155/2014/563165>
- Pereira, E.G. & Nascimento, L.B. (2004) Descrição da vocalização e do girino de *Pseudopaludicola mineira* Lobo, 1994, com notas sobre a morfologia de adultos (Amphibia, Anura, Leptodactylidae). *Arquivos do Museu Nacional*, 62, 233–240.
- R Development Core Team (2012) *The R Project for Statistical Computing*. Vienna, Austria. Available from: <http://www.r-project.org/> (accessed 30 January 2013)
- Roberto, I.J., Cardozo, D. & Ávila, R.W. (2013) A new species of *Pseudopaludicola* (Anura, Leiuperidae) from western Piauí state, northeast Brazil. *Zootaxa*, 3636 (2), 348–360.
<http://dx.doi.org/10.11646/zootaxa.3636.2.6>
- Santos, T.G., Rossa-Feres, D.C. & Casatti, L. (2007) Diversidade e distribuição espaço-temporal de anuros em região com pronunciada estação seca no sudeste do Brasil. *Iheringia*, 97, 37–49.
<http://dx.doi.org/10.1590/S0073-47212007000100007>
- Silva, R.A., Martins, I.A. & Rossa-Feres, D.C. (2008) Bioacústica e sítio de vocalização em taxocenoses de anfíbios de áreas aberta no noroeste paulista. *Biota Neotropica*, 8, 123–134.
- Silva, R.A., Martins, I.A. & Rossa-Feres, D.C. (2011) Environmental heterogeneity: Anuran diversity in homogeneous environments. *Zoologia*, 28, 610–618.
<http://dx.doi.org/10.1590/S1984-46702011000500009>
- Silva, F.R., Candreira, C.P. & Rossa-Feres, D.C. (2012) Dependence of anuran diversity on environmental descriptors in farmland ponds. *Biodiversity and Conservation*, 21, 1411–1424.
<http://dx.doi.org/10.1007/s10531-012-0252-z>
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011) MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Molecular Biology and Evolution*, 28, 2731–2739.
<http://dx.doi.org/10.1093/molbev/msr121>
- Taylor, W.R. & Van Dyke, G.C. (1985) Revised procedures for staining and clearing small fishes and other vertebrates for bone and cartilage study. *Cybium*, 9, 107–119.
- Toledo, L.F., Siqueira, S., Duarte, T.C., Veiga-Menoncello, A.C.P., Recco-Pimentel, S.M. & Haddad, C.F.B. (2010) Description of a new species of *Pseudopaludicola* Miranda-Ribeiro, 1926 from the state of São Paulo, Southeastern Brazil (Anura, Leiuperidae). *Zootaxa*, 2498, 38–48.
- Veiga-Menoncello, A.C.P., Lourenço, L.B., Strüssmann, C., Rossa-Feres, D.C., Andrade, G.V., Giaretta, A.A. & Recco-Pimente, S.M. (2014) A phylogenetic analysis of *Pseudopaludicola* (Anura) providing evidence of progressive chromosome reduction. *Zoologica Scripta*, 43, 261–272.
<http://dx.doi.org/10.1111/zsc.12048>
- Vences, M., Thomas, M., Bonett, R.M. & Vieites, D.R. (2005) Deciphering amphibian diversity through DNA barcoding: chances and challenges. *Philosophical Transactions of the Royal Society Biological Sciences*, 360, 1859–1868.
<http://dx.doi.org/10.1098/rstb.2005.1717>
- Vences, M., Köhler, J., Vieites, D.R. & Glaw, F. (2011) Molecular and bioacoustic differentiation of deep conspecific lineages of the Malagasy treefrogs *Boophis tampoka* and *B. luteus*. *Herpetology Notes*, 4, 239–246.
- Vieites, D.R., Wollenberg, K.C., Andreone, F., Köhler, J., Glaw, F. & Vences, M. (2009) Vast underestimation of Madagascar's biodiversity evidenced by an integrative amphibian inventory. *Proceedings of the National Academy of Sciences of the U.S.A.*, 106, 8267–8272.

APPENDIX 1. Voucher specimens examined.

Pseudopaludicola facureae: BRAZIL: MINAS GERAIS: **Uberlândia**: Males: AAG-UFU 853–855; 2277–8; 2528; 2622; 2681; 3586; 3588; UFMT 17745–5; 17748; and ZUEC 13650–2. Females: AAG-UFU 2279; 2282; 3585; 3587; 3589; 3591; 4731 and ZUEC 13653–4.

APPENDIX 2. Analyzed sound files.

Pseudopaludicola atragula sp. nov.: BRAZIL: SÃO PAULO: **Itém**: LH 702 (voucher specimen UFMT 16198); LH 703 (voucher specimen UFMT 16206); LH 439–440 (no voucher specimens).

Pseudopaludicola facureae: BRAZIL: MINAS GERAIS: **Uberlândia**: LH 721–723 (no voucher specimens); LH 724 (voucher specimen UFMT 17745); LH 725 (voucher specimen UFMT 17746); LH 726 (voucher specimen UFMT 17748).