

<http://dx.doi.org/10.11646/zootaxa.3941.2.3>
<http://zoobank.org/urn:lsid:zoobank.org:pub:2550E24F-6A14-4FE4-A505-CB0C832EB7BB>

Description of *Pseudingolfiella possessionis* n. sp. (Crustacea, Amphipoda) from sub-Antarctic Île de La Possession, Crozet archipelago: the second freshwater amphipod known from the Antarctic biome, a human introduction of Gondwanan ancestry?

WILLEM H. DE SMET

Department of Biology, University of Antwerp Campus Drie Eiken, Universiteitsplein 1, B-2610 Wilrijk, Belgium.
E-mail: willem.desmet@uantwerpen.be

Abstract

A new species of freshwater amphipod, *Pseudingolfiella possessionis* n. sp. (Senticaudata, Pseudingolfiellidae), is described from the submerged moss vegetation of small brooklets at sub-Antarctic Île de La Possession, Crozet archipelago. It constitutes the second freshwater amphipod species known for the sub-Antarctic and Antarctic region, and the fourth member of the genus. The main characters distinguishing it from all congeners are: the spine on the posterior margin of the dactylus, incisor and lamina mobilis of mandible each with 5 teeth, the setation of the maxilliped, the vestigial second article of pleopod 3 in the female, the undulate and laterally notched posterolateral margin of the external ramus of uropods 1 and 2 in the male, the spinulate dorsomedian projection of the telson.

Key words: Antarctica, diversity, *Pseudingolfiella*, biogeography, chorology

Introduction

About 1,870 amphipod species and subspecies are recognized from fresh or inland waters world-wide, accounting for 20% of the total known amphipod diversity (Väinölä *et al.* 2008). The described systematic diversity of freshwater amphipods from the sub-Antarctic and Antarctic (Pugh *et al.* 2002; Väinölä *et al.* 2008) is confined to a single species, *Kergueleniola macra* (Ruffo, 1970), family Kergueleniolidae Lowry & Myers, 2013, that has so far only been reported once. It was described by Ruffo (1970, 1974) on a single specimen found in the stomach of a freshwater trout, *Salvelinus fontinalis* (Mitchill, 1814), from the sub-Antarctic Îles Kerguelen. An unidentified amphipod species was mentioned from benthic vegetation of South Georgia by Dartnall & Heywood (1980). During a study on freshwater micro-invertebrates of the sub-Antarctic island Île de La Possession, Crozet archipelago (Îles Crozet), a new species of amphipod belonging to the genus *Pseudingolfiella* was encountered, which is described here.

Material and methods

The Crozet archipelago ($45^{\circ}57'–46^{\circ}27'S$, $50^{\circ}15'–52^{\circ}15'E$) consists of 5 small volcanic islands situated in the Southern Indian Ocean (Fig. 1), just north of the Antarctic Convergence. Biogeographically it belongs to the cold temperate province of Stonehouse (1982). The main island Île de La Possession ($46^{\circ}25'S$ – $51^{\circ}45'E$) has a surface area of 147 km^2 , and is characterized by an oceanic and cold climate. The mean annual temperature is 5.1°C ; little variation is recorded over the year, with a mean minimum temperature of 3°C and mean maximum temperature of 7.9°C (Météo-France, 1970–94 records). The annual precipitation usually surpasses 2,200 mm, with 293 rain-days per year. Strong winds, which may exceed 200 km h^{-1} and usually coming from SW-NNW, scourge the island throughout the year.

References

- Bousfield, E.L. (1982) Amphipoda. In: Parker, S.J. (Ed.), *Synopsis and classification of living organisms*. Vol 2. McGraw-Hill Book Comp, New York, pp. 254–294.
- Bousfield, E.L. (1989) Revised morphological relationships within the amphipod genera *Pontoporeia* and *Gammaracanthus* and the ‘Glacial Relict’ significance of their postglacial distributions. *Canadian Journal of Fisheries and Aquatic Sciences*, 46, 1714–725.
- Boxshall, G.A. & Jaume, D. (2009) Exopodites, epipodites and gills in crustaceans. *Arthropod Systematics & Phylogeny*, 67, 229–254.
- Chevallier, L., Nougier, J. & Cantagrel, J.M. (1982) Volcanology of Possession Island, Crozet archipelago (T.A.A.F.). In: Craddock, C. (Ed.), *Antarctic geosciences. IUGS, B, 4*, University of Wisconsin Press, Madison, pp. 652–658.
- Coineau, N. (1977) Le genre *Pseudedingolfiella* aux Kerguelen. *CNFRA*, 42, 287–294.
- Crispo, E. (2007) The Baldwin effect and genetic assimilation: revisiting two mechanisms of evolutionary change mediated by phenotypic plasticity. *Evolution*, 61, 2469–2479.
<http://dx.doi.org/10.1111/j.1558-5646.2007.00203.x>
- Dartnall, H.J.G. & Heywood, R.B. (1980) The freshwater fauna of South Georgia. *British Antarctic Survey Bulletin*, 50, 115118.
- de Bovée, F., Coineau, N., Soyer, J. & Travé, J. (1973) Sur l’existence d’une faunule interstitielle littorale dans l’Archipel de Kerguelen (Terres Australes et Antarctiques françaises). *Comptes Rendus des Séances de l’Académie des Sciences, Paris, Séries D* (276), 1041–1044.
- Delépine, G. (1995) *Les Iles Australes Françaises, Kerguelen, Crozet, Amsterdam, Saint-Paul*. Editions Ouest-France, Rennes, 213 pp.
- Derycke, S., Remerie, T., Backeljau, T., Vierstraete, A., Vanfleteren, J., Vincx, M. & Moens, T. (2008) Phylogeography of the *Rhabditis (Pellioidoditis) marina* species complex: evidence for long-distance dispersal, and for range expansions and restricted gene flow in the northeast Atlantic. *Molecular Ecology*, 17, 3306–3322.
- De Smet, W.H. (2003) A new record of *Limnognathia maerski* Kristensen & Funch, 2000 (Micrognathozoa) from the subantarctic Crozet Islands, with redescription of the trophi. *Journal of Zoology, London*, 258, 381–393.
- Giret, A. (1987) Géologie des Terres Australes Françaises. *CNFRA*, 58, 17–41.
- Grosso, L.E., Peralta, M.A. & Ruffo, S. (2006) Description of *Pseudoingolfiella morimotoi*, sp. nov. (Crustacea, Amphipoda) from New Zealand and transantarctic distribution of the genus. *Subterranean Biology*, 4, 67–77.
<http://dx.doi.org/10.3354/meps025169>
- Hay, W.H., DeConto, R.M., Wold, C.N., Wilson, K.M., Voigt, S., Schulz, M., Wold, A.R., Dullo, W.-C., Ronov, A.B., Balukhovsky, A.N. & Söding, E. (1999) Alternative global Cretaceous paleogeography. In: Barrera, E. & Johnson, C.C. (Eds.), *Evolution of the Cretaceous Ocean-Climate System*. Boulder, Colorado, Geological Society of America, Special Paper, 332, 1–47.
- Highsmith, R.C. (1985) Floating and algal rafting as potential dispersal mechanisms in brooding invertebrates. *Marine Ecology Progress Series* 25, 169–179.
<http://dx.doi.org/10.3354/meps025169>
- Kikuchi, S., Matsumasa, M. & Yashima, Y. (1993) The ultrastructure of the sternal gills forming a striking contrast with the coxal gills in a fresh-water amphipod (Crustacea). *Tissue and Cell*, 25, 915–928.
- Kikuchi, S. & Matsumasa, M. (1997) Ultrastructural evidence for osmoregulatory function of the sterna epithelia in some gammaridean amphipods. *Journal of Crustacean Biology*, 17, 377–388.
- Kristensen, R.M. & Funch, P. (2000) Micrognathozoa: a new class with complicated jaws like those of Rotifera and Gnathostomulida. *Journal of Morphology*, 246, 1–49.
- Lincoln, R.J. & Hurley, D.E. (1981) The calceolus, a sensory structure of gammaridean amphipods (Amphipoda: Gammaridea). *Bulletin of the British Museum of Natural History (Zoology)*, 40, 103–116.
- Lowry, J.K. & Myers, A.A. (2012) New, mainly southern hemisphere, freshwater families of Amphipoda (Crustacea), together with a description of the first freshwater calliopiid, *Lutriwita bradburyi* gen. nov. et sp. nov. *Zootaxa*, 3499, 27–45.
- Lowry, J.K. & Myers, A.A. (2013) A phylogeny and classification of the Senticaudata subord. nov. (Crustacea: Amphipoda). *Zootaxa*, 3610 (1), 1–80.
<http://dx.doi.org/10.11646/zootaxa.3610.1.1>
- Mitchill, S.L. (1814) Report, in part, of Samuel L. Mitchill, M. D., ... on the fishes of New-York. New York, pp. 1–28.
- Noodt, W. (1959) Estudios sobre crustáceos chilenos de aguas subterráneas. I. *Ingolfiella chilensis* n. sp. de la playa marina de Chile Central. (Crustacea, Amphipoda). *Investigaciones Zoológicas Chilenas*, 5, 199–209.
- Noodt, W. (1965) Interstítiole Amphipoden der konvergenten Gattungen *Ingolfiella* Hansen und *Pseudedingolfiella* n. gen. aus Südamerika. *Crustaceana*, 9, 17–30.
- Ochyra, R. (1998) *The moss flora of King George Island, Antarctica*, Szafer Institute of Botany, PAS, Cracow, XXIV, 279 pp.
- Pugh, P.J.A., Dartnall, H.J.G. & McInnes, S.J. (2002) The non-marine Crustacea of Antarctica and the islands of the Southern Ocean: biodiversity and biogeography. *Journal of Natural History*, 36, 1047–1103.
- Ruffo, S. (1970) Descrizione di *Kerguelenella macra* n. gen. n. sp. (Amphipoda Gammaridae) delle isole Kerguelen. *Memorie del Museo civico di Storia naturale di Verona*, 18, 43–54.
- Ruffo, S. (1974) Note brevi: Due cambiamenti di nome di generi di crostacei anfipodi. *Bollettino del Museo civico di Storia*

- naturale di Verona*, 1, 1–507.
- Schalke, H.J.W. & Van Zinderen Bakker, E.M. (1971) History of the vegetation. In: Van Zinderen Bakker, E.M., Winterbottom, J.M. & Dyer, R.A. (Eds.), *Marion and Prince Edward Islands, report on the South African biological and geological expedition 1965–1966*. A.A. Balkema, Cape Town, pp. 88–97.
- Soyer, J. (1976) Recherches sur la faune mésopsammique dans l'archipel de Kerguelen. *Comité national français des recherches antarctiques*, 39, 107–141.
- Stonehouse, B. (1982) La zonation écologique sous les hautes latitudes australes. *Comité national français des recherches antarctiques*, 51, 531–537.
- Väinölä, R., Witt, J.D.S., Grabowski, M., Bradbury, J.H., Jazdzewski, K. & Sket, B. (2008) Global diversity of amphipods (Amphipoda; Crustacea) in freshwater. *Hydrobiologia*, 595, 241–255.
- Van der Putten, N., Verbruggen, C., Ochyra, R., Verleyen, E. & Frenot, Y. (2010) Subantarctic flowering plants: pre-glacial survivors or post-glacial immigrants? *Journal of Biogeography*, 37, 582–592.
- Watling, L. (1989) A classification system for crustacean setae based on the homology concept. In: Felgenhauer, B.E., Watling, L. & Thistle, A.B. (Eds.), *Functional Morphology of Feeding and Grooming in Crustacea*. A.A. Balkema, Rotterdam, Brookfield. *Crustacean Issues*, 6, 15–26.
- Whitman, D. & Agrawal, A.A. (2005) What is phenotypic plasticity and why is it important? In: Ananthakrishnan, T.N. & Whitman, D.W. (Eds.), *Insect Phenotypic Plasticity: Diversity of Responses*. Science Publishers, Enfield, pp. 1–63.