



<http://dx.doi.org/10.11646/zootaxa.3949.1.9>

<http://zoobank.org/urn:lsid:zoobank.org:pub:F05517F9-B4ED-417C-AAA4-6FBA488D8C0F>

First report of *Toumeyella parvicornis* (Cockerell) in Europe (Hemiptera: Coccidae)

ANTONIO P. GARONNA, SALVATORE SCARPATO, FRANCESCO VICINANZA & BRUNO ESPINOSA

Laboratorio di Entomologia “E. Tremblay”, Dipartimento di Agraria, Università degli Studi di Napoli “Federico II”, Via Università, 100, 80055 Portici, Napoli, Italy. E-mail: garonna@unina.it

Toumeyella parvicornis (Cockerell), the Pine tortoise scale, is reported for the first time from Europe. Pre-imaginal instars and adults were observed starting from the fall of 2014 in several municipalities of Campania (Italy). Since the species is recorded as a harmful pest of several pine species in North America and the Caribbean, the presence of this alien insect in Europe is alarming and represents a possible major threat for native pine species and related economic activities.

Scale insects are one of the most commonly transported groups of insects in plant trade and at the same time are one of the most successful invasive groups of insects (Pellizzari *et al.*, 2005; Malumphy *et al.*, 2012). Here we report the occurrence of *Toumeyella parvicornis* (Cockerell), the Pine tortoise scale, in southern Italy infesting *Pinus pinea* L. in urban environments. Italy is one of the most invaded of European countries by alien insects (Inghilesi *et al.*, 2013) and this new record updates the list of the established Coccoidea in the country to 50 species (Mazzeo *et al.*, 2014).

During the last months of 2014, severe infestations of a sap feeding insect, producing huge amount of honeydew with impressive sooty mould development, were observed on stone pines (*P. pinea*) in Naples and neighboring municipalities (Campania, Italy). Abundant colonies of a soft scale insect were collected in many sampling sites. All slide mounted specimens (i.e., adult females and males, male tests and first-instar nymphs), representative of the collected samples, were ascribed to a species of the genus *Toumeyella* Cockerell associated with Pinaceae (Hamon & Williams, 1984; Hodgson, 1994; Kondo & Pellizzari, 2011; Miller & Williams, 1995; Sheffer & Williams, 1990), namely *T. parvicornis*. The genus *Toumeyella* has a wide distribution in the Nearctic and Neotropical Regions, with most species being described from the USA (Williams & Kondo, 2008). Currently there are 18 named species included in this genus, which are distributed in Brazil (3 spp.), Canada (2 spp.), Colombia (1 sp.), Cuba (2 spp.), Guatemala (1 sp.), Mexico (6 spp.), Puerto Rico (1 sp.), Turks & Caicos Islands (1 sp.), the USA (9 spp.) and Venezuela (1 sp.) (Kondo & González, 2014; Ben-Dov *et al.*, 2015).

Four species of *Toumeyella* live on *Pinus* spp.: *T. parvicornis*, *T. pini* (King), the striped pine scale, *T. pinicola* Ferris, the irregular pine scale, and *T. virginiana* Williams & Kosztarab, the Virginia pine scale (Williams & Kondo, 2008). The Pine tortoise scale has two forms, the typical bark form and a leaf form (Hamon & Williams 1984; Kondo & Pellizzari, 2011) (Fig. 1a–c). This species can be easily separated from other congeneric species by the characteristic dorsal bilocular pore clusters present on the adult female, with other minor differences on the antennae, anal plates and spiracular furrows (Fig. 2a–c) (Hamon & Williams, 1984; Williams & Kosztarab, 1972). *Toumeyella* species can be also distinguished by morphological characters of the first instars (Figs 1d, 2d) and male tests (Miller & Williams, 1990; Sheffer & Williams, 1990). *T. parvicornis* was originally described from Florida (USA), on *Pinus palustris* and *P. taeda* but is now known from Canada, Mexico, and all of the USA and is an important pest of several *Pinus* species (Clarke, 2013). Recently, *T. parvicornis* has been accidentally introduced into Puerto Rico (Segarra-Carmona & Cabrera-Asencio, 2010) and Turks and Caicos Islands, where it has become a severe pest of *Pinus caribaea* (Malumphy *et al.*, 2012). *T. parvicornis* shows great environmental adaptability and an easy acceptance of new hosts, and this might allow the species to spread quickly into new territories. It is considered a destructive species infesting the most vigorously growing succulent twigs. In the northern parts of its range, the pine tortoise scale has one generation per year whereas four generations are typical for southern Georgia, USA (Clarke, 2013). In Turks and Caicos Islands, where even a fifth generation is possible, the impact of *T. parvicornis* has been severe. As reported by Malumphy *et al.* (2012), some of the infested areas contain no live trees or seedlings whilst other areas have a few live pines amongst dead and moribund trees. The rapid population growth rate of pine tortoise scale in Caribbean pine forests has been attributed to a combination of warm climatic conditions and a lack of natural enemies (Malumphy *et al.*, 2012).




FIGURE 3. *Toumeyella parvicornis* (Cockerell), distribution map in Italy (Campania). Municipalities listed in alphabetic order with latitude and longitude coordinates of sample points [Acerra (1), 40°56'03"N, 14°21'59"E; Afragola (2), 40°54'46"N, 14°17'57"E; Arzano (3), 40°55'00"N, 14°16'28"E; Caivano (4), 40°57'59"N, 14°18'11"E; Casoria (5), 40°53'33"N, 14°19'06"E; Cercola (6), 40°51'12"N, 14°21'11"E; Ercolano (7), 40°49'12"N, 14°22'41"E; Giugliano in Campania (8), 40°55'43"N, 14°02'19"E; Marcianise (9), 41°00'17"N, 14°19'20"E; Napoli (10–20), 40°49'34"N, 14°11'18"E; 40°47'53"N, 14°10'47"E; 40°50'06"N, 14°08'48"E; 40°50'25"N, 14°09'34"E; 40°49'12"N, 14°12'11"E; 40°48'52"N, 14°11'58"E; 40°50'12"N, 14°18'44"E; 40°50'12"N, 14°18'51"E; 40°51'50"N, 14°15'47"E; 40°52'28"N, 14°14'18"E; 40°51'53"N, 14°13'36"E; Nola (21), 40°57'30"N, 14°27'46"E; Portici (22), 40°48'42"N, 14°20'37"E; San Giorgio a Cremano (23–24), 40°49'46"N, 14°20'19"E; 40°49'32"N, 14°20'48"E; San Sebastiano al Vesuvio (25), 40°50'12"N, 14°22'07"E; Torre del Greco (26), 40°47'52"N, 14°22'57"E].

References

- Ben-Dov, Y., Miller, D.R. & Gibson, G.A.P. (2014) ScaleNet: a database of the scale insects of the world. Available from: <http://www.sel.barc.usda.gov/scalenet/scalenet.htm> (accessed 1 January 2015)
- Bracalini, M., Benedettelli, S., Croci, F., Terreni, P., Tiberi, R. & Panzavolta, T. (2013) Cone and seed pests of *Pinus pinea*: assessment and characterization of damage. *Journal of Economic Entomology*, 106 (1), 229–234. <http://dx.doi.org/10.1603/EC12293>
- Clarke, S.R. (2013) Pine Tortoise Scale. USDA Forest Service. *Forest Insect & Disease Leaflet*, 57, 1–8.
- Cristinzio, G., Bosso, L., Somma, S., Varlese, R. & Saracino A. (2014) *Serious damage by Diplodia africana on Pinus pinea in the Vesuvius National Park (Campania Region, southern Italy)*. II International Congress of Silviculture, Florence, Italy, November 26–29, 2014, poster 2.1.
- Hamon, A.B. & Williams, M.L. (1984) *The Soft Scale Insects of Florida (Homoptera: Coccoidea: Coccidae). Arthropods of Florida and Neighboring Land Areas. Vol. 11*. Florida Department of Agriculture and Consumer Service Division of Plant Industry, Gainesville, 194 pp.
- Hodgson, C.J. (1994) *The Scale Insect Family Coccidae: an identification manual to genera*. CAB International,

Wallingford, Oxon, 639 pp. [UK]

- Inghilesi, A.F., Mazza, G., Cervo, R., Gherardi, F., Sposimo, P., Tricarico, E. & Zapparoli, M. (2013) Alien insects in Italy: Comparing patterns from the regional to European level. *Journal of Insect Science*, 13 (73). Available from: <http://jinsectscience.oxfordjournals.org/content/jis/13/1/73.full.pdf> (accessed 15 April 2015)
- Kondo, T. & González, H. (2014) A new species of *Toumeyella* Cockerell (Hemiptera: Coccidae) on *Myrtillocactus geometrizans* (Cactaceae) from Mexico with a checklist of known species of *Toumeyella*. *Insecta Mundi*, 0396, 1–10.
- Kondo, T. & Pellizzari, G. (2011) Description of a new species of *Toumeyella* Cockerell (Hemiptera: Coccidae) from Mexico; with a taxonomic key to Mexican species. *Revista Brasileira de Entomologia*, 55 (2), 229–233. <http://dx.doi.org/10.1590/S0085-56262011005000019>
- Malumphy, C., Hamilton, M.A., Manco, B.N., Green, P.W.C., Sanchez, M.D., Corcoran, M. & Salamanca, E. (2012) *Toumeyella parvicornis* (Hemiptera: Coccidae) causing severe decline of *Pinus caribaea* var. *bahamensis* in the Turks and Caicos Islands. *Florida Entomologist*, 95, 113–119. <http://dx.doi.org/10.1653/024.095.0118>
- Mazzeo, G., Longo, S., Pellizzari, G., Porcelli, F., Suma, P. & Russo, A. (2014) Exotic Scale Insects (Coccoidea) on Ornamental Plants in Italy: a Never-Ending Story. *Acta zoologica bulgarica*, 6 (Supplement), 55–61.
- Miller, G.L. & Williams, M.L. (1995) Systematic analysis of the adult males of *Toumeyella* group, including *Mesolecanium nigrofasciatum*, *Neolecanium cornuparvum*, *Pseudophilippia quaintancii* and *Toumeyella* spp. (Homoptera: Coccidae) from America north of Mexico. *Contributions of the American Entomological Institute*, 28, 1–68.
- Miller, G.L. & Williams, M.L. (1990) Tests of male soft scale insects (Homoptera: Coccidae) from America north of Mexico, including a key to the species. *Systematic Entomology*, 15 (3), 339–358. <http://dx.doi.org/10.1111/j.1365-3113.1990.tb00070.x>
- Pellizzari, G., Dalla Montà, L. & Vacante, V. (2005) Alien insect and mite pests introduced to Italy in sixty years (1945–2004). Plant Protection and Plant Health in Europe: Introduction and Spread of Invasive Species. *Proceedings BCPC Symposium*, 81, 275–276.
- Segarra-Carmona, A.E. & Cabrera-Asencio, I. (2010) *Toumeyella parvicornis* (Cockerell) (Hemiptera: Coccoidea: Coccidae): a new invasive pest of pine trees in Puerto Rico. *Journal of the Agricultural University of Puerto Rico*, 94, 175–177.
- Sheffer, B.J. & Williams, M.L. (1990) Descriptions, distribution, and host-plant records of eight first instars in the genus *Toumeyella* (Homoptera: Coccidae). *Proceedings of the Entomological Society of Washington*, 92 (1), 44–57.
- Williams, M.L. & Kondo, T. (2008) Status and current composition of the soft scale insect genus *Toumeyella* (Hemiptera: Coccoidea). *Proceedings of the XIth International Symposium on Scale Insect Studies*. September 24–27, 2007, 29–32. [Branco, M., Franco, J.C. & Hodgson, C. (Eds), Oeiras, Portugal]
- Williams, M.L. & Kosztarab, M. (1972) Morphology and systematics of the Coccidae of Virginia with notes on their biology (Homoptera: Coccoidea). *Virginia Polytechnic Institute and State University, Research Division Bulletin*, 74, 1–215.