

ZOOTAXA

4015

A taxonomic revision of the *Phrynosoma douglasii* species complex (Squamata: Phrynosomatidae)

RICHARD R. MONTANUCCI¹

¹*Department of Biological Sciences, Clemson University, Clemson, SC 29634-0314.*
E-mail: RRMNT@clemson.edu



Magnolia Press
Auckland, New Zealand

RICHARD R. MONTANUCCI

A taxonomic revision of the *Phrynosoma douglasii* species complex (Squamata: Phrynosomatidae)
(Zootaxa 4015)

177 pp.; 30 cm.

11 Sept. 2015

ISBN 978-1-77557-789-8 (paperback)

ISBN 978-1-77557-790-4 (Online edition)

FIRST PUBLISHED IN 2015 BY

Magnolia Press

P.O. Box 41-383

Auckland 1346

New Zealand

e-mail: zootaxa@mapress.com

<http://www.mapress.com/zootaxa/>

© 2015 Magnolia Press

All rights reserved.

No part of this publication may be reproduced, stored, transmitted or disseminated, in any form, or by any means, without prior written permission from the publisher, to whom all requests to reproduce copyright material should be directed in writing.

This authorization does not extend to any other kind of copying, by any means, in any form, and for any purpose other than private research use.

ISSN 1175-5326 (Print edition)

ISSN 1175-5334 (Online edition)

Table of contents

Abstract	3
Introduction	3
Materials and methods	5
Results and discussion	10
Taxonomic accounts	25
<i>Phrynosoma brevirostris</i> Girard 1858a	25
<i>Phrynosoma bauri</i> sp. nov.	33
<i>Phrynosoma diminutum</i> sp. nov.	39
<i>Phrynosoma douglasii</i> (Bell 1828)	45
<i>Phrynosoma hernandesi</i> <i>hernandesi</i> Girard 1858a	51
<i>Phrynosoma hernandesi</i> <i>ornatum</i> Girard 1858b, comb. nov.	61
<i>Phrynosoma ornatissimum</i> <i>ornatissimum</i> Girard 1858a	67
<i>Phrynosoma ornatissimum</i> <i>brachycercum</i> Smith 1942, comb. nov.	75
Hybridization between <i>Phrynosoma hernandesi</i> and <i>Phrynosoma ornatissimum</i>	82
Summary	110
Acknowledgements	111
References	112
Appendix I	117
Appendix II	163

Abstract

Short-horned lizards (*Phrynosoma douglasii* species complex) occur throughout the inter-montane West and Great Plains of western North America. The comparative morphology and color pattern variation of short-horned lizards was studied in 3,174 specimens. Multivariate analyses of 20 morphological and color-pattern characters were applied to 977 specimens, and univariate statistics were summarized for 52 samples totaling 1,134 specimens. The results of the morphological data analyses support the recognition of *P. douglasii* (Bell 1828) as a distinct species, and the resurrection of *P. brevirostris* Girard 1858a and *P. ornatissimum* Girard 1858a as species distinct from *P. hernandesi* Girard 1858a. Two new species allied to *P. brevirostris* are described: *P. bauri* sp. nov. from the eastern plains of Colorado and northeastern New Mexico, southeastern Wyoming and southwestern Nebraska south of the North Platte River, and *P. diminutum* sp. nov. endemic to the San Luis Valley of southern Colorado and northern New Mexico. The Mexican taxon *brachycercum* Smith 1942 is reassigned as a subspecies of *P. ornatissimum*, based on non-discrete character differences and overall morphometric similarity. The ranges of *P. hernandesi* and *P. ornatissimum* broadly overlap in central New Mexico, the former taxon occupying the coniferous forests of disjunct mountain ranges, the latter occurring in the surrounding desert grasslands. Principal components analysis has revealed morphological evidence of hybridization where the two taxa meet, generally within ecotones between montane forest associations and grasslands. Principal components analysis has also revealed a high level of morphological variability in populations occurring in the Colorado Plateau region of northeastern Arizona, northwestern New Mexico, extreme southwestern Colorado and adjacent Utah. The evidence suggests that these populations arose through past hybridization between the two species. The taxon *ornatum* Girard 1858b, although sharing several traits with *P. brevirostris*, is morphologically close to *P. hernandesi*. It is regarded as a stabilized population of hybrid origin, but treated taxonomically as a subspecies of *P. hernandesi*. The taxonomic arrangement in this study, with the exception of *P. douglasii*, is largely discordant with the proposed taxonomy from a previously published study based on mitochondrial DNA sequence data.

Key words: geographic variation, morphology, *Phrynosoma*, Phrynosomatidae, species delimitation, systematics, taxonomy

Introduction

As presently conceptualized, the *Phrynosoma douglasii* species complex consists of two species, *Phrynosoma douglasii* (Bell) and *Phrynosoma hernandesi* Girard, following Zamudio *et al.* (1997). The former species occurs along the Cascade Range in the Pacific Northwest region and the deserts as far east as southern Idaho (Stebbins, 1985; Nussbaum *et al.* 1983). The latter is widely distributed from the mixed grass prairies of southern Alberta and southwestern Saskatchewan (Powell & Russell, 1998) southward through the woodlands and grasslands of the Great Basin, Rocky Mountains, western Great Plains and eastern slopes of the Sierra Madre Occidental, reaching central Zacatecas, Mexico (Stebbins, 1985; Smith & Flores-Villela, 1994).