



Erratum

<http://dx.doi.org/10.11646/zootaxa.3999.4.10>

<http://zoobank.org/urn:lsid:zoobank.org:pub:93984950-A3A8-4F57-84AB-997D3C54DB16>

AWAL RIYANTO, L. LEE GRISMER & PERRY L. WOOD, JR. (2015) *Cyrtodactylus rosichonariefi* sp. nov. (Squamata: Gekkonidae), a new swamp-dwelling bent-toed gecko from Bunguran Island (Great Natuna), Indonesia. *Zootaxa*, 3964 (1): 114–124.

Abstract should read as follows:

Cyrtodactylus rosichonariefi sp. nov. is the second newly described species of Bent-toed gecko from Bunguran Island (Great Natuna), Indonesia. This species occurs in the Sekunyam Forest and is differentiated from all other species of the *C. semenanjungensis* species group of the Thai-Malay Peninsula by having the following unique combination of characteristics: intermediate size (SVL of adult male holotype 54.6 mm); enlarged femoral scales, no femoral pores, no precloacal groove, enlarged precloacal scales, no precloacal pores; subcaudals not enlarged; and dorsal pattern blotched but lacking paired, dark, semilunar-shaped blotches on upper nape. The new species is the sister species of *C. majulah* Grismer, Wood & Lim.