

<http://dx.doi.org/10.11646/zootaxa.4060.1.16>  
<http://zoobank.org/urn:lsid:zoobank.org:pub:E57EEA21-F6D3-4032-B7B4-8FAEAD74D574>

## Annotated checklist of eels (orders Anguilliformes and Saccopharyngiformes) from Taiwan

HSUAN-CHING HO<sup>1,2</sup>, DAVID G. SMITH<sup>3</sup>, JOHN E. MCCOSKER<sup>4</sup>, YUSUKE HIBINO<sup>5</sup>, KAR-HOE LOH<sup>6</sup>, KENNETH A. TIGHE<sup>3</sup> & KWANG-TSAO SHAO<sup>7,\*</sup>

<sup>1</sup>National Museum of Marine Biology & Aquarium, Pingtung, Taiwan

<sup>2</sup>Institute of Marine Biology, National Dong Hwa University, Pingtung, Taiwan

<sup>3</sup>Smithsonian Institution, Museum Support Center, Suitland, MD, U.S.A.

<sup>4</sup>California Academy of Sciences, San Francisco, U.S.A.

<sup>5</sup>Fisheries Research Laboratory, Mie University, Mie, Japan

<sup>6</sup>Institute of Ocean and Earth Sciences, University of Malaya, Malaysia

<sup>7</sup>Biodiversity Research Center, Academia Sinica, Taipei, Taiwan

\*Corresponding author. E-mail:zoskt@gate.sinica.edu.tw

### Abstract

An annotated checklist of eels, orders Anguilliformes and Saccopharyngiformes, occurring in Taiwanese waters is presented. The checklist is the result of a series of systematic studies conducted by the authors in the past few years. The eel fauna of Taiwan is one of the richest in the world with a total of 206 species in 74 genera and 13 families in Anguilliformes and a single species in Saccopharyngiformes. The most species-rich families are the Muraenidae with 71 species, followed by the Ophichthidae with 60 species, the Congridae with 29 species, and the Synaphobranchidae with 17 species. Moreover, three genera and 42 species have been described based on at least one type specimen collected from Taiwan. Of these, 36 species are recognized as valid and 23 species are known only from Taiwanese waters at present. Historical records of all Taiwanese eel species are reviewed by examining the original descriptions and figures, vouchers, as well as the recently collected specimens, where available. This represents the first detailed checklist of eels from Taiwanese waters.

**Key words:** Taxonomy, elopomorpha, Anguilliformes, Saccopharyngiformes, Taiwan

### Introduction

Taiwan has a rich and diverse fish fauna. The island is located on the Tropic of Cancer, resulting in a unique mixture of tropical and temperate species. The fishes of Taiwan have been studied over the years by many ichthyologists, both foreign and domestic. The history of ichthyology in Taiwan was summarized by Ho & Shao (2011), and that of taxonomy of anguilliform eels is summarized in the present special issue (Ho *et al.*, 2015a).

The first assessment of the eel fauna of Taiwan was compiled by Chen & Weng (1967), who recognized 10 families with 65 species, and included four newly described species. Chen & Yu (1986) increased the number to 87 species and in a comprehensive book "*Fishes of Taiwan*", Shen *et al.* (1993) recognized 9 families and 108 species. In the past two decades, 17 eel species were described from Taiwan (Ho & Shao, 2011; Ho *et al.*, 2015a). The Taiwan Fish Database (Shao, 2015) currently lists 13 families with 168 species of eels. However, many taxonomic problems remain and many groups need further study. Among these groups are the eels of the order Anguilliformes.

During the course of their systematic studies on the eels of Taiwan, the contributors to this volume examined many specimens previously collected and deposited in fish collections in Taiwan and elsewhere. They also made new collections from around the island and exchanged information on the taxonomic results. As a result, many new species were discovered and many previously described species were recorded from Taiwan for the first time (McCosker *et al.*, 2009, 2012; Hibino *et al.*, 2012; Ho *et al.*, 2013); more are waiting to be described.

The main purpose of present work is to provide a detailed checklist that includes all known species in Taiwan and their current status and to provide a solid foundation for future studies.

## Methods and materials

The checklist is arranged alphabetically by family, then by genera and by species. The new species described from Taiwan are marked \* (including those in the synonymies) and species newly recorded in this volume are marked +. Voucher specimens are provided if a new record is not mentioned in other sources.

Following the scientific name, a common name in Chinese and ‘local’ synonymies are provided for all Taiwanese taxa, unless otherwise indicated. The first author who reported a particular species from Taiwan is underlined in the synonymies.

Current status follows the studies made by the authors in the present issue or those listed in *Catalog of Fishes* (Eschmeyer, 2015), unless otherwise indicated. A Remarks section includes ecological notes, field observations, taxonomic history, nomenclature or systematics is provided, if applicable. The new genera and species described in this volume are presented here in the checklist. Several of the species groups are still under investigation and their taxonomic status may change in the future.

Institutional acronyms follow Eschmeyer (2015). The geographic coordinates for localities can be found in Ebert *et al.* (2013:table 5, fig. 1).

## Results

Two hundred and six species in 13 families and 74 genera of the order Anguilliformes and one species in one family and one genus of the order Saccopharyngiformes are recognized in this work, which gathers together all information from previous references and our recent study results. Of these, 71 species belong to the Muraenidae, 60 to the Ophichthidae, 29 to the Congridae and 17 to the Synaphobranchidae which represent the most species rich families in Taiwan waters.

Three new genera and 42 species have been described from Taiwan, of which 36 species are valid and 23 of them are currently only found in Taiwan. There are also 1 family and 29 new species records added to the Taiwanese fauna. Moreover, we recognized at least nine undescribed species which are not included in this checklist.

### Annotated checklist of eels of Taiwan

#### Order Anguilliformes

鰻形目

#### Family Anguillidae

鰻科

##### *Anguilla bicolor pacifica* Schmidt, 1928

太平洋雙色鰻

*Anguilla pacifica* Schmidt, 1928:190, 206 (Type locality: Pacific, north of the equator).

*Anguilla bicolor pacifica* Schmidt, 1928: Tseng & Tabeta, 1983:28; Chen & Yu, 1986:242; Lin, 2007:86; Chen, 2009:4; Chou & Kao, 2011:33; Chang *et al.*, 2011:1; Chang *et al.*, 2013:44.

*Anguilla bicolor* McClelland, 1844: Shen, 1984a:103.

Remarks. Chen (2009) notes that this species is mainly collected from fresh waters of eastern Taiwan.

### ***Anguilla celebesensis* Kaup, 1856**

西里伯斯鰻

*Anguilla celebesensis* Kaup, 1856b:42 (type locality: Lake Tondano, Sulawesi, Indonesia). Tseng, 1982:59; Chang *et al.*, 2011:1; Chen *et al.*, 2012:1; Chang *et al.*, 2013:44.

Remarks. Tseng (1982) first recorded this species based on juveniles. Although there are other records of this species, no adults have been formally recorded. Chen *et al.* (2012) noted that some juveniles of this species may have been included with *Anguilla luzonensis*, a recently described species. Chang *et al.* (2013) confirmed that this species, also based on juveniles, is present in Taiwanese waters.

### ***Anguilla japonica* Temminck & Schlegel, 1846**

日本鰻

*Anguilla japonica* Temminck & Schlegel, 1846:258 (type locality: Japan). Oshima, 1919:263; Chen, 1956:73; Liaw, 1960:4; Chen & Weng, 1967:5; Chen, 1969:124; Shen, 1984a:103; Chen & Yu, 1986:242; Shen *et al.*, 1993:97, fig. 14-5; Shao *et al.*, 1994a:273; Chen & Fang, 1999:51; Chen, 2004:26; Lin, 2007:82; Chen, 2009:6; Ho *et al.*, 2010:table 1; Chou & Kao, 2011:34; Ho & Shao, 2011:21, table 1; Chang *et al.*, 2011:1; Chang *et al.*, 2013:44.

*Anguilla bostoniensis* (not of Lesueur, 1817):?Günther, 1870:31.

\**Anguilla remifera* Jordan & Evermann, 1902:325, fig. 7 (type locality: Taiwan). Jordan & Richardson, 1909:171, fig. 8.

*Anguilla sinensis* McClelland, 1843: Chen, 1956:73.

*Muraena japonica* (Temminck & Schlegel, 1846): Fowler, 1932a:49, fig. 1.

Remarks. Günther (1870) recorded *Anguilla bostoniensis* based on several half-grown specimens collected from Taiwan, but their identities are not verified since whereabouts of these specimens are unknown.

### ***Anguilla luzonensis* Watanabe, Aoyama & Tsukamoto, 2009**

呂宋鰻

*Anguilla luzonensis* Watanabe, Aoyama & Tsukamoto, 2009:389 (type locality: north Luzon Island, Philippines). Chen *et al.*, 2012:68; Chang *et al.*, 2013:44.

Remarks. Chen *et al.* (2012) noted that adults of this species were collected from Lan-yu Island and juveniles were collected in northeastern and southwestern Taiwan. Chang *et al.* (2013) reported that juveniles occur in Taiwan, although no adults were collected.

### ***Anguilla marmorata* Quoy & Gaimard, 1824**

鱸鰻

*Anguilla marmorata* Quoy & Gaimard, 1824:241 (type locality: Waigeo, Indonesia). Chen, 1956:73; Chen & Weng, 1967:4; Chen, 1969:124; Shen, 1984a:103; Chen & Yu, 1986:242; Shen *et al.*, 1993:97, fig. 14-6; Chen & Fang, 1999:52; Lin, 2007:84; Chen, 2009:8; Ho *et al.*, 2010:table 1; Chou & Kao, 2011:35; Chang *et al.*, 2011:1.

*Anguilla mauritiana* Bennett, 1831: Jordan & Evermann, 1902:325; Oshima, 1919:262.

*Anguilla manillensis* (Bleeker, 1864a): Jordan & Richardson, 1909:171.

Remarks. This is the most widely distributed species of *Anguilla*, occurring from East Africa to French Polynesia, and from Papua New Guinea to southern Japan.

## **Family Chlopsidae**

草鰻科

### **\**Chlopsis nanhaiensis* Tighe, Ho, Pogonoski & Hibino, 2015**

南海草鰻

*Chlopsis nanhaiensis* Tighe, Ho, Pogonoski & Hibino in Tighe et al., 2015:22, figs. 4–7 (type locality: Dong-gang, southwestern Taiwan).

Remarks. This is a rare species known from the two type specimens collected by bottom trawl off southwestern Taiwan.

#### ***Chilorhinus platyrhynchus* (Norman, 1922)**

扁吻草鰻

*Brachyconger platyrhynchus* Norman, 1922:218 (type locality: New Britain Island, Bismarck Archipelago [Papua New Guinea, western Pacific]).

*Chilorhinus platyrhynchus* (Norman, 1922): Shao et al., 2008:237.

Remarks. This is a rare species represented by a few specimens collected by bottom trawl from southwestern Taiwan and two catalog records (three specimens total) in the USNM collection collected by rotenone off southern Taiwan.

#### **+*Kaupichthys japonicus* Matsubara & Asano, 1959**

日本眶鼻鰻

*Kaupichthys diodontus japonicus* Matsubara & Asano, 1959:293, figs. 1, 2A–B (type locality: Near Kametsu, Tokunoshima, Amami Islands, Japan).

*Kaupichthys hyoprorooides* (not of Strömmann, 1896): Shao et al., 2008:237.

*Kaupichthys diodontus* (not of Schultz, 1943): Chen & Yu, 1986:255.

Voucher specimens: ASIZP 54931 (Heng-chun, Pingtung). Also represented by ten catalog records (41 specimens total) in the USNM collection collected by rotenone off southern Taiwan.

Remarks. Previous Taiwanese records of *K. hyoprorooides* and *K. diodontus* represent misidentifications of *K. japonicus* and are not valid. The only voucher present in Taiwan is reidentified as the present species. It is notable that there are four rows (two on each side) of blunt teeth on the vomer, whereas the original description described only two, one on each side. However, one of us (KAT) has observed a number of specimens of *Kaupichthys* that have two rows of teeth on each side of the vomer.

### **Family Colocongridae**

短康吉鰻科

#### ***Coloconger japonicus* Machida, 1984**

日本短康吉鰻

*Coloconger japonicus* Machida in Okamura & Kitajima, 1984:87 (type locality: Okinawa Trough, 750–760 meters). Shao et al., 2008:239.

Remarks. This is a rare species represented by only two specimens collected in southwestern Taiwan.

#### **+*Coloconger raniceps* Alcock, 1889**

蛙頭短康吉鰻

*Coloconger raniceps* Alcock, 1889:456 (Type locality: Andaman Sea off Ross Island).

Voucher specimen: ASIZP 63790 (SW Taiwan).

Remarks. This is a new record for Taiwan and it is represented by a single specimen collected in southwestern Taiwan at 821 m.

## **Family Congridae**

康吉鰻科

### ***Ariosoma anago* (Temminck & Schlegel, 1846)**

白錐體糯鰻

*Conger anago* Temminck & Schlegel, 1846:259 (type locality: Nagasaki, Japan).

*Leptocephalus anagoides* (not of Bleeker, 1854): Jordan & Richardson, 1909:171.

*Alloconger anagoides* (not of Bleeker, 1854): Chu, 1957:14; Lee & Yang, 1966:54, fig. 2; ?Chen & Weng, 1967:45; Chen, 1969:132; Chen & Yu, 1986:252; Shen et al., 1993:115; Shao et al., 1994a:274.

*Ariosoma anago* (Temminck & Schlegel, 1846): Shao et al., 2008:239.

Remarks. There has been considerable confusion over this name in the past. The species treated as *Anago* [=*Ariosoma*] *anago* by Asano (1962) is in fact *Ariosoma meeki* (Jordan & Snyder, 1901); the true *A. anago* was treated by Asano as *Ariosoma anagoides* (Bleeker, 1852). Most subsequent authors followed Asano. Previous records of *Ariosoma anago* from Taiwan are reidentified by us as *A. meeki*, and records of *A. anagoides* are reidentified as *A. anago*. Known from Japan and Taiwan; southern limit of distribution uncertain.

### ***Ariosoma fasciatum* (Günther, 1872)**

條紋錐體糯鰻

*Poeciloconger fasciatus* Günther, 1872a:673 (type locality: Manado, Sulawesi, Indonesia).

\**Ariosoma nancyae* Shen, 1998:10, figs. 3A-B, 5C, 6C (type locality: Nan-fang-ao fish market, northeastern Taiwan).

*Ariosoma fasciatum* (Günther, 1872a): Ho & Shao, 2011:23, table 1.

Remarks. This is a rare species collected by hook and line off eastern Taiwan. The head pore pattern is not typical of *Ariosoma* and further investigation is needed to determine its generic status.

### ***Ariosoma major* (Asano, 1958)**

大錐體糯鰻

*Alloconger shiroanago major* Asano, 1958:191 (off Shibushi, Kagoshima Prefecture, Japan). Lee & Yang, 1966:55, fig. 3.

*Alloconger major* Asano, 1958: Chen & Weng, 1967:47; Chen, 1969:132; Shen 1984a:112; Chen & Yu, 1986:252; Shao et al., 2008:239.

*Ariosoma shiroanago major* (Asano, 1958): Shen, 1998:10.

*Ariosoma major* (Asano, 1958): Ho et al., 2010:table 1.

Remarks. This is a common species collected by bottom trawl around Taiwan.

### ***Ariosoma meeki* (Jordan & Snyder, 1900)**

米氏錐體糯鰻

*Congrellus meeki* Jordan & Snyder, 1900:347 (type locality: Tokyo Bay, Japan).

*Anago anago* (non Temminck & Schlegel, 1846): Chu, 1957:15; Lee & Yang, 1966:53, fig. 1; Chen & Weng, 1967:43; Lee, 1967:44; Chen, 1969:132; Shen 1984a:112; Chen & Yu, 1986:252; Shen, 1988:15; Shao et al., 1994a:274.

*Ariosoma anago* (non Temminck & Schlegel, 1846): Chen, 1956:76; Liang & Yeh, 1964:8; Shen, 1998:8.

*Ariosoma meeki* (Jordan & Snyder, 1900): Chen, 2004:31; Shao et al., 2008:239.

Remarks. This species was referred to *Anago anago* in most previous works. It is the most abundant *Ariosoma* species in bottom trawl catches in Taiwan.

### **+*Ariosoma shiroanago* (Asano, 1958)**

銀錐體糯鰻

*Alloconger shiroanago shiroanago* Asano, 1958:193 (type locality: off Owashi, Mie Prefecture, Japan).

Voucher specimen: NMMB-P17984 (Dong-gang).

Remarks. This is a new record for Taiwan. It is rare and only represented by a few specimens from Taiwan.

**+*Bathycongrus guttulatus* (Günther, 1887)**

小斑深海糯鰻

*Congromuraena guttulata* Günther, 1887:252 (type locality: off Matuku, Fiji Islands).

Voucher specimens: USNM 401083 (Dong-gang); NMMB-P 4041 (Dong-gang).

Remarks. This is a new record for Taiwan. It is a rare species collected by bottom trawl off southwestern Taiwan.

***Bathycongrus retrotinctus* (Jordan & Snyder, 1901)**

網格深海糯鰻

*Leptocephalus retrotinctus* Jordan & Snyder, 1901:853 (type locality: Tokyo fish market, Japan).

*Congrina retrotincta* (Jordan & Snyder, 1901): Lee & Yang, 1966:59, fig. 8; Chen & Weng, 1967:55; Chen, 1969:132; Shen 1984a:113; Chen & Yu, 1986:253.

*Bathycongrus retrotinctus* (Jordan & Snyder, 1901): Shao *et al.*, 2008:239.

Remarks. This is a common species collected by bottom trawl off northeastern and southwestern Taiwan, though not as abundant as *B. wallacei*.

**+*Bathycongrus wallacei* (Castle, 1968)**

瓦氏深海糯鰻

*Congrina wallacei* Castle, 1968:709 (type locality: off the mouth of Limpopo River, Mozambique, 480–500 m).

Voucher specimens: Many specimens in ASIZP and NMMB-P.

Remarks. Previously recorded from Japan and the western Indian Ocean (Castle & Smith, 1999), this is a new record for Taiwan. It is a very common species collected by bottom trawl off northeastern and southwestern Taiwan.

***Bathymyrus simus* Smith, 1965**

銼吻淵油鰻

*Bathymyrus simus* Smith, 1965:8, fig. 1B, pl. 1 (type locality: Vietnam). Chen & Weng, 1967:43; Chen, 1969:130; Shen 1984a:111; Chen & Yu, 1986:252; Shen *et al.*, 1993:116; Shao *et al.*, 2008:239; Chen, 2004:31; Ho *et al.*, 2010:table 1.

Remarks. This is a rare species only found off southwestern Taiwan.

**+*Bathyuroconger parvibranchialis* (Fowler, 1934)**

小鰓深海尾糯鰻

*Silvesterina parvibranchialis* Fowler, 1934:275, fig. 35 (type locality: Butung [Buton] Strait, Philippines, 5°35'00"S, 122°20'00"E, 559 fathoms [1022 m]).

*Bathyuroconger parvibranchialis* (Fowler, 1934): Smith, 1999b:1686.

Voucher specimens: ASIZP 63789 (SW Taiwan); ASIZP 63794 (SW Taiwan); ASIZP 66181 (SW Taiwan).

Remarks. Recent studies by DGS indicate that the type material of this species actually contains two species. Further studies are needed to determine which of these occurs in Taiwan. They are characterized by their small gill opening. Also reported from Japan (F. Tashiro, pers. comm.).

***Bathyuroconger vicinus* (Vaillant, 1888)**

深海尾糯鰻

*Uroconger vicinus* Vaillant, 1888:86, pl. 6, fig. 1 (Coasts of Soudan, Arguin Bank, and Cape Verde Islands, 932–1633 m).  
*Bathyuroconger vicinus* (Vaillant, 1888): Shao *et al.*, 2008:239.

Remarks. As currently recognized, this species occurs worldwide. It is collected occasionally from deep waters by bottom trawl off northeastern Taiwan.

***Blachea xenobranchialis* Karrer & Smith, 1980**

奇鰓鰻

*Blachea xenobranchialis* Karrer & Smith, 1980:643, figs. 1–6 (type locality: Mozambique Channel, 12°43'S, 48°15'E, 348–360 m). Ho & Shao, 2010:286.

Remarks. This is a rare species represented by one specimen from Taiwan and a few additional specimens from elsewhere in the Indo-Pacific.

***Conger cinereus* Rüppell, 1871**

灰糯鰻

*Conger cinereus* Rüppell in Klunzinger 1871:607 (type locality: Red Sea). Chen & Weng, 1967:50; Chen, 1969:132; Shen 1984a:112; Chen & Yu, 1986:252; Shen *et al.*, 1993:116; Shao *et al.*, 2008:239.

*Conger wilsoni* (not of Bloch & Schneider, 1801): ?Chen, 2004:32.

Remarks. This is a common species in coral reef habitats.

**+*Conger jordani* Kanazawa, 1958**

喬氏糯鰻

*Conger jordani* Kanazawa, 1958:250, fig. 4 (type locality: Misaki, Kanagawa Prefecture, Japan)

Voucher specimens: ASIZP 62593 (Daxi).

Remarks. This is a common species in Taiwan. Previous records of *C. japonicus* may refer to this species.

**+*Conger macrocephalus* Kanazawa, 1958**

大頭糯鰻

*Conger macrocephalus* Kanazawa, 1958:254, pl. 1, fig. M (Verde Island passage, Luzon Island, Philippines, 13°34'37"N, 121°07'30"E, ca. 180 fathoms [329 m]).

*Conger japonicus* (not of Bleeker, 1879): ?Chen & Weng, 1967:49; ?Chen, 1969: 132; ?Shen 1984a:112; ?Shen, 1988:15; ?Shen *et al.*, 1993:116; ?Chen, 2004:32; ?Ho *et al.*, 2010:table 1.

*Conger wilsoni* (not of Bloch & Schneider, 1801): ?Chen & Yu, 1986:252.

*Conger conger* (not of Linnaeus, 1758): ?Lee & Yang, 1966:56, fig. 5.

Voucher specimens: Many in ASIZP, NMMP-P and USNM.

Remarks. This is a new record for Taiwan. This species appears to be common in deep waters around Taiwan and is mainly collected by bottom trawl. Many specimens in collections were previously mis-identified as *C. japonicus*.

***Conger myriaster* (Brevoort, 1856)**

繁星糯鰻

*Anguilla myriaster* Brevoort, 1856:282, pl. 11, fig. 2 (type locality: Hakodate, Hokkaido, Japan).

*Conger myriaster* (Brevoort, 1856): Lee & Yang, 1966:56, fig. 4; Chen & Weng, 1967:48; Chen, 1969: 132; Shen 1984a:112; Chen & Yu, 1986:252; Shen *et al.*, 1993:116; Shao *et al.*, 2008:239; Ho *et al.*, 2010:table 1.  
*Conger japonicus* Bleeker, 1879:32, pl. 2, fig. 2 (type locality: Japan). Kanazawa, 1958:244).

Remarks. This species is commonly collected by bottom trawl around Taiwan. The identity of *Conger japonicus* has been disputed for many years, but current studies (Smith *et al.*, in prep.) indicate that Kanazawa (1958) was correct in synonymizing it with *C. myriaster*. The identity of many references to *Conger japonicus* in Taiwan is uncertain.

### ***Congriscus megastomus* (Günther, 1877)**

大口糯鰻

*Congromuraena megastoma* Günther, 1877:445 (type locality: off Inoshima, Hiroshima Prefecture, Japan).  
*Congriscus megastomus* (Günther, 1877): Ho *et al.*, 2010:table 1.

Remarks. This species is occasionally collected by bottom trawl off northeastern Taiwan.

### **+*Gnathophis asanoi* Karmovskaya, 2004**

淺野氏領吻鰻

*Gnathophis asanoi* Karmovskaya, 2004:S18, fig. 10 (type locality: Philippines, Sibuyan Sea, 13°08'N, 122°39'E, 440–280 m).

Voucher specimen: USNM 401080 (Dong-gang).

Remarks. This species originally described from the Philippines and newly recorded in Taiwan.

### ***Gnathophis heterognathos* (Bleeker, 1858)**

異領領吻糯鰻

*Myrophis heterognathos* Bleeker, 1858b:9, pl. 3, fig. 1 (type locality: Nagasaki, Japan).  
*Gnathophis heterognathos* (Bleeker, 1858): Kaup, 1859:7; Castle, 1963:17.

*Rhynchocymba nystromi* (Jordan & Snyder, 1901): Chen & Weng, 1967:51; Chen, 1969:132; Shen 1984a:112; Chen & Yu, 1986:252.

*Rhynchocymba nystromi nystromi* (Jordan & Snyder, 1901): Chu, 1957:14; Liang & Yeh, 1964:8; Lee & Yang, 1966:57, fig. 6;  
Shao *et al.*, 2008:239.

*Gnathophis nystromi* (Jordan & Snyder, 1901): Shen, 1988:15; Shao *et al.*, 1994a:274; Chen, 2004:32.

*Gnathophis nystromi ginanago* (Asano, 1958): ?Shao *et al.*, 1994a:274.

Remarks. This species was referred to *Gnathophis nystromi* in most previous works. However, Castle (1963) first suggested and Karmovskaya (2004) later confirmed that *G. nystromi* is a junior synonym of *G. heterognathos*, which has not previously been used for the Taiwanese specimens.

### ***Gorgasia japonica* Abe, Miki & Asai, 1977**

日本園鰻

*Gorgasia japonica* Abe, Miki & Asai, 1977:1, pl. 1, fig. 2 (type locality: near Hachijo-Kojima, 33°07'N, 139°42'E, Japan).  
Shao, 1990:11.

Remarks. This is a rare species and has only been recorded from Lan-yu Island off southeastern Taiwan.

### **\**Gorgasia taiwanensis* Shao, 1990**

臺灣園鰻

*Gorgasia taiwanensis* Shao, 1990:4, figs.2-6 (Type locality: Wanlitung, southern Taiwan, 22 m). Shao *et al.*, 2008:239; Ho & Shao, 2011:23, table 1.

Remarks. This is a rare species occasionally found on sandy bottoms near coral reef areas in southern and southeastern Taiwan.

***Heteroconger hassi* (Klausewitz & Eibl-Eibesfeldt, 1959)**

哈氏異糯鰻

*Xarifania hassi* Klausewitz & Eibl-Eibesfeldt, 1959:138, figs. 6, 8, 15, 16 (type locality: Addu Atoll, between Gan Island and Fedu Island, Maldives Islands, Indian Ocean).

*Heteroconger hassi* (Klausewitz & Eibl-Eibesfeldt, 1959): Shao *et al.*, 2008:239.

Remarks. This species is commonly found in colonies with up to several hundred individuals, on sandy bottom near coral reef areas around Taiwan.

***Japononconger sivicolus* (Matsubara & Ochiai, 1951)**

禿吻日本糯鰻

*Arisoma* [sic] *sivicola* Matsubara & Ochiai, 1951:11 (type locality: off Owase, Kumano-Nada, Japan).

*Japononconger sivicolus* (Matsubara & Ochiai, 1951): Shao *et al.*, 2008:239.

Remarks. A rare species only represented by several specimens collected from deep waters around Taiwan.

**+*Macrocephenchelys brachialis* Fowler, 1934**

大頭糯鰻

*Macrocephenchelys brachialis* Fowler, 1934:277, fig. 36 (type locality: Makassar Strait, between Borneo and Sulawesi, Indonesia, 2°56'00"S, 118°47'30"E, 367 fathoms [671 m]).

Voucher specimens: many specimens in ASIZP, NMMB-P and USNM.

Remarks. This is a new record for Taiwan. It is commonly collected from deep waters off northeastern and southwestern Taiwan. It is quite abundant in the bycatch of bottom trawls.

**\**Macrocephenchelys brevirostris* (Chen & Weng, 1967)**

大頭短吻糯鰻

*Rhynchoconger brevirostris* Chen & Weng, 1967:54, fig. 40 (type locality: Tungkang, southwestern Taiwan). Chen, 1969:132;

Shen 1984a:113; Chen & Yu, 1986:253; Shen *et al.*, 1993:117.

*Macrocephenchelys brevirostris* (Chen & Weng, 1967): Smith in Randall & Lim, 2000:586; Karmovskaya, 2004:S24; Shao *et al.*, 2008:239; Ho *et al.* 2010:23, table 1; Ho & Shao, 2011:24, table 1.

Remarks. This is a common deepwater species collected by bottom trawls, more abundant off southwestern than in northeastern Taiwan. It is likely a senior synonym of *Macrocephenchelys soela* Castle, 1990, described from Australia. According to Karmovskaya (2004), this species also occurs in the Philippines and New Caledonia.

**+*Parabathymyrus brachyrhynchus* (Fowler, 1934)**

粗頷擬海糯鰻

*Arisoma* [sic] *brachyrhynchus* Fowler, 1934:269, fig. 30 (Bongo Island, Illana Bay, southern Mindanao Island, Philippines, 7°21'45"N, 124°07'15"E, 158 fathoms [289 m]).

*Parabathymyrus brachyrhynchus* (Fowler, 1934): Smith & Kanazawa, 1977:532; Ho *et al.*, 2015b:134.

Remarks. A common species collected from deep waters by bottom trawl off northeastern and southwestern Taiwan, more abundant in southwestern Taiwan. Many specimens in the collections were previously misidentified as *P. macrophthalmus*.

### ***Parabathymyrus macrophthalmus* Kamohara, 1938**

大眼擬海鰻鰻

*Parabathymyrus macrophthalmus* Kamohara, 1938:14, fig. 4 (type locality: Mimase, Kochi Prefecture, Japan). Li, 1960:87; Chen & Weng, 1967:41; Chen, 1969:132; Shen 1984a:111; Chen & Yu, 1986:252; Shen *et al.*, 1993:117; Shao *et al.*, 2008:239; Ho *et al.*, 2015b:137.

Remarks. This species is less abundant than *P. brachyrhynchus* in Taiwan. We also examined many specimens collected from Vietnam, which represent a new record in Vietnamese waters.

### **\**Rhynchoconger ectenurus* (Jordan & Snyder, 1901)**

突吻纖尾鰻鰻

*Leptocephalus ectenurus* Jordan & Richardson, 1909:171, pl. 66 (type locality: Takao [Kaohsiung], southwestern Taiwan).

*Rhynchoconger ectenurus* (Jordan & Richardson, 1909): Chen, 1956:76; Lee & Yang, 1966:58, fig. 7; Chen & Weng, 1967:52; Chen, 1969:132; Shen 1984a:113; Chen & Yu, 1986:253; Shen *et al.*, 1993:117; Shao *et al.*, 1994a:274; Shao *et al.*, 2008:239; Ho *et al.*, 2010:table 1; Ho & Shao, 2011:23, table 1.

*Rhynchocymba ectenura* (Jordan & Snyder, 1901): Chu, 1957:15.

\**Rhynchoconger brachuata* Chu & Chen, 1958:127, fig. 1 (type locality: Taipei fish market).

Remarks. A very common species collected by bottom trawls from deep waters around Taiwan.

### ***Uroconger lepturus* (Richardson, 1845)**

狹尾鰻鰻

*Congrus lepturus* Richardson, 1845b:106, pl. 56, figs. 1–6 (type locality: Canton, China).

*Uroconger lepturus* (Richardson, 1845): Chen & Weng, 1967:56; Chen, 1969:133; Shen 1984a:113; Chen & Yu, 1986:253; Shen *et al.*, 1993:118; Shao *et al.*, 2008:239; Ho *et al.*, 2010:table 1.

Remarks. A species complex which may represent more than one species. At least two species occur in Taiwan, based on DNA barcodes, number of cephalic pores and coloration. Further studies are needed. Very common around Taiwan.

## **Family Derichthyidae**

項鰻科

### **+*Nessorhamphus danae* Schmidt, 1931**

丹氏鴨項鰻

*Nessorhamphus danae* Schmidt, 1931:487, fig. 4 (type locality: Indian Ocean, W of Sumatra).

Voucher specimen: USNM 436661.

Remarks. This is a new record for the family and genus in Taiwan and is based on a single metamorphic larva from Dong-gang, SW Taiwan. It is tentatively identified as *N. danae* based on the position of the last vertical blood vessel (LVBV, or renal artery), at approximately myomere 65–71, which is within the range given for this species by Karmovskaya (1985). The comparable count in *N. ingolfianus*, the only other species in the genus, is 72–78.

## **Family Moringuidae**

鰐鰻科

Remarks. The Moringuidae is undoubtedly the most misunderstood family of eels and one of the most misunderstood of all fish families. It needs to be thoroughly revised. Three species have been recorded in Taiwan,

but these records should be considered provisional. *Moringua macrochir* was recorded by Chou & Kao (2011). However, no voucher was provided and the identification is thus not verified.

### ***Moringua abbreviata* (Bleeker, 1863)**

短線鰐鰻

*Aphthalmichthys abbreviatus* Bleeker, 1863:163 (type locality: Prigi, Java; Batu; Manado, Sulawesi; Ternata; Ambon, Indonesia).

*Moringua abbreviata* (Bleeker, 1863): Chen & Weng, 1967:8; Chen, 1969:124; Shen, 1984a:103; Chen & Yu, 1986:247; Shen et al., 1993:98, fig. 14-8; Shao et al., 2008:237; Ho et al., 2010:table 1.

### ***Moringua macrocephalus* (Bleeker, 1863)**

大頭鰐鰻

*Aphthalmichthys macrocephalus* Bleeker, 1863:165 (type locality: Timor Island, southern Malay Archipelago): Li, 1960:88.

*Moringua macrocephalus* (Bleeker, 1863): Chen & Weng, 1967:7; Chen, 1969:124; Shen, 1984a:103; Chen & Yu, 1986:242; Shen et al., 1993:98, fig. 14-9; Shao et al., 2008:237; Ho et al., 2010:table 1.

## **Family Muraenesocidae**

海鰻科

### ***Congrhyynchus talabonoides* Fowler, 1934**

原鶴康吉吻鰻

*Congrhyynchus talabonoides* Fowler, 1934:273, fig. 33 (type locality: Macabalan Pt., northern Mindanao Island, Philippines, 8°37'37"N, 124°35'E, ca. 214 fathoms [391 m]). Chen & Yu, 1986:254.

Remarks. This species has only been recorded in Taiwan once; there is a voucher in NMMB-P collection (NMMB-P1461) and the identification is confirmed.

### ***Congresox talabon* (Cuvier, 1829)**

原鶴海鰻

*Conger talabon* Cuvier, 1829:350 (type locality: Vizagapatnam, India).

*Muraenesox talabon* (Cuvier, 1829): Li, 1960:84; Chen & Weng, 1967:77; Chen, 1969:133; Shen 1984a:114; Chen & Yu, 1986:254; Shen, 1988:15; Chen, 2004:31.

*Congresox talabon* (Cuvier, 1829): Castle & Williamson, 1975:3.

Remarks. This species was first reported from Taiwan by Li (1960), but the specimens on which the record was based were reidentified by Lin et al. (2013) as *Oxyconger leptognathus*. Chen & Weng (1967) provided an account of the species from Taiwan and stated that specimens were "...preserved in the Department of Zoology of the National Taiwan University and in the Taiwan Fisheries Research Institute, collected from Keelung & Suao, length 500–1000 mm," although they did not give catalog numbers. Lin et al. (2013), however, were apparently unable to find these specimens and stated that "there was not any real record of *C. talabon* in Taiwan." Castle & Williamson (1975: fig. 4) included Taiwan in the range of *Congresox talabonoides* (Bleeker, 1853a) but not *C. talabon*. Further studies are needed to confirm the presence of either of these species in Taiwan.

### **\**Gavialiceps taiwanensis* (Chen & Weng, 1967)**

台灣鱸頭鰻

*Chlopsis taiwanensis* Chen & Weng, 1967:81, fig. 61 (Type locality: Tungkang, southwestern Taiwan). Chen, 1969:134; Shen, 1984a:114; Chen & Yu, 1986:255; Shen et al., 1993:115.

*Gavialiceps taiwanensis* (Chen & Weng, 1967): Karmovskaya, 1994:84; Smith in Randall & Lim, 2000:586; Shao et al., 2008:239; Ho et al., 2010:25; Ho & Shao, 2011:24, table 1; Lin et al., 2013:167.

*Gavialiceps taeniola* (not of Alcock, 1889): Shao *et al.*, 2008:239.

Remarks. The assignment of this species to the Muraenesocidae is provisional. This is the most abundant eel species found in bycatch of midwater trawls in Taiwan.

### ***Muraenesox bagio* (Hamilton, 1822)**

百吉海鰻

*Muraena bagio* Hamilton, 1822:24, 364 (type locality: Ganges River estuaries, India).

*Muraenesox bagio* (Hamilton, 1822): Shen, 1984a:114; Chen & Yu, 1986:254; Shen *et al.*, 1993:118; Chen & Fang, 1999:55; Shao *et al.*, 2008:239; Chen, 2004:31; Shen & Wu, 2011:135; Lin *et al.*, 2013:169.

*Muraenesox yamaguchiensis* Katayama & Takai, 1954:97, Figs. 1–2 (type locality: Yamaguchi Prefecture, Japan). Chen & Weng, 1967:76; Chen, 1969:133.

Remarks. *Muraenesox yamaguchiensis* was synonymized with *M. bagio* by Castle & Williamson (1975). This is a very common species collected by bottom trawls and hook and lines around Taiwan.

### ***Muraenesox cinereus* (Forsskål, 1775)**

灰海鰻

*Muraena cinerea* Forsskål, 1775:22 (Type locality: Jeddah, Saudi Arabia, Red Sea).

*Muraenesox cinereus* (Forsskål, 1775): Chen, 1956:77; Chu, 1957:15; Chen & Weng, 1967:75; Chen, 1969:133; Shen 1984a:114; Chen & Yu, 1986:254; Shen *et al.*, 1993:118; Shao *et al.*, 1994a:276; Chen, 2004:31; Shao *et al.*, 2008:239; Ho *et al.*, 2010:table 1; Shen and Wu, 2011:135; Lin *et al.*, 2013:170.

Remarks. This is a very common species collected by bottom trawls and hook and lines around Taiwan.

### ***Oxyconger leptognathus* (Bleeker, 1857)**

狹頷海鰻

*Conger leptognathus* Bleeker, 1857c:27, pl. 2, fig. 2 (type locality: Nagasaki, Japan).

*Oxyconger leptognathus* (Bleeker, 1857): Li, 1960:850; Chen & Weng, 1967:79; Chen, 1969:133; Shen 1984a:113; Chen & Yu, 1986:254; Shen, 1988:15; Shen *et al.*, 1993:119; Chen, 2004:31; Shao *et al.*, 2008:239; Ho *et al.*, 2010:table 1; Lin *et al.*, 2013:171.

Remarks. An uncommon species collected by bottom trawls from around Taiwan. Based on a recent field observation by HH, this species may live in association with deep corals in southwestern Taiwan at depths around 100–200 m.

## **Family Muraenidae**

鯉科

### ***Anarchias allardicei* Jordan & Starks, 1906**

褐裸臀鯉

*Anarchias allardicei* Jordan & Starks in Jordan & Seale, 1906:204, fig. 9 (type locality: Pago Pago, Tutuila Island, American Samoa). Lee, 1980:57; Shen *et al.*, 1993:99, fig. 19-2; Chen *et al.*, 1994:54; Chen, 2004:26; Shao *et al.*, 2008:238; Loh *et al.*, in press:table 1.

*Gymnomuraena concolor* (not of Rüppell, 1838): Chen & Weng, 1967:38; Chen, 1969:129; Shen, 1984a:107; Chen & Yu, 1986:249.

Remarks. A small species commonly collected in tidal pools or subtidal zone of southern Taiwan.

***Anarchias cantonensis* (Schultz, 1943)**

廣東裸臀鯽

*Uropterygius cantonensis* Schultz, 1943:27, pl. 4, fig. 3a (type locality: lagoon of Canton Island, Phoenix Islands, southern central Pacific).

*Anarchias cantonensis* (Schultz, 1943): Loh *et al.*, in press:table 1.

Remarks. A rare species collected in tidal pools or subtidal zone in southeastern Taiwan.

***Channomuraena vittata* (Richardson, 1845)**

環帶裂口鯽

*Ichthyophis vittatus* Richardson, 1845a: pl. 53, figs. 7–9 (type locality uncertain, brought from China).

*Channomuraena vittata* (Richardson, 1845): Loh *et al.*, in press:table 1.

Remarks. A very rare species only represented by a few specimens collected from southeastern Taiwan.

**\**Cirrimaxilla* Chen & Shao, 1995**

*Cirrimaxilla* Chen & Shao, 1995:328 (Type species. *Cirrimaxilla formosa* Chen & Shao, 1995, by monotypy). Loh *et al.*, 2008:138; Séret *et al.*, 2008:192; Ho & Shao, 2011:21.

**\**Cirrimaxilla formosa* Chen & Shao, 1995**

台灣鬚鯽

*Cirrimaxilla formosa* Chen & Shao, 1995:330, fig. 1 (Type locality: Nanwan, southern tip of Taiwan, Ping-tung County). Shao *et al.*, 2008:238; Ho & Shao, 2011:21; tale 1; Loh *et al.*, 2008:138; Loh *et al.*, in press:table 1.

Remarks. This species was tentatively synonymized with *Channomuraena bauchotae* from the Indian Ocean by Böhlke & Smith (2002), but this needs to be confirmed. At any rate, neither of these species belongs in *Channomuraena*, which has quite different dentition. *Cirrimaxilla formosa* was recently found to be common in stomach contents of sea snakes in New Caledonia (Séret *et al.*, 2008).

***Echidna nebulosa* (Ahl, 1789)**

星帶蝮鯽

*Muraena nebulosa* Ahl, 1789:7, pl. 1, right fig. (type locality: in Oriental India [Indonesia]).

*Echidna nebulosa* (Ahl, 1789): Chen & Weng, 1967:13; Chen, 1969:127; Shen, 1984a:106; Shen, 1988:11; Chen & Yu, 1986:247; Shen *et al.*, 1993:100; Shao *et al.*, 1994a:276; Chen *et al.*, 1994:47; Chen *et al.*, 1997:appendix 1; Chen, 2004:27; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

Remarks. A very common species found in coral reef regions of shallow waters around Taiwan.

***Echidna polyzona* (Richardson, 1845)**

多環蝮鯽

*Muraena polyzona* Richardson, 1845b:112, pl. 55, figs. 11–14 (No locality stated).

*Echidna polyzona* (Richardson, 1845): Jordan & Richardson, 1909:173; Chen, 1956:79; Chen & Weng, 1967:11; Chen, 1969:127; Shen, 1984a:106; Shen, 1988:11; Chen & Yu, 1986:247; Shen *et al.*, 1993:100, fig. 15-1, 2; Shao *et al.*, 1994a:276; Chen *et al.*, 1994:47; Chen *et al.*, 1997:appendix 1; Chen, 2004:27; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

Remarks. A common species found in the coral reef regions around Taiwan.

***Echidna xanthospilos* (Bleeker, 1859)**

黃斑蝮鯀

*Muraena xanthospilos* Bleeker, 1859:348 (type locality: Java, Indonesia, eastern Indian Ocean).  
*Echidna xanthospilos* (Bleeker, 1859): Loh *et al.*, in press:table 1.

Remarks. A rare and relatively deepwater species living in southeastern Taiwan that is represented by only a few specimens in collections.

***Enchelycore bayeri* (Schultz, 1953)**

貝氏勾吻鯀

*Gymnothorax bayeri* Schultz in Schultz *et al.*, 1953:124, figs. 23f, 26 (type locality: lagoon coral head at Kieschiechi Island, Rongelap Atoll, Marshall Islands, western Pacific, 20 feet [6 m]).  
*Enchelycore bayeri* (Schultz, 1953): Loh *et al.*, in press:table 1.

Remarks. This is a rare species in Taiwan that is only represented by a few specimens in the collections.

***Enchelycore bikiniensis* (Schultz, 1953)**

比吉尼勾吻鯀

*Gymnothorax bikiniensis* Schultz in Schultz *et al.*, 1953:116, figs. 23e, 24 (Type locality: Cherry Island, Bikini Atoll, Marshall Islands).  
*Enchelycore bikiniensis* (Schultz, 1953): Shen *et al.*, 1993:100, fig. 15-3; Chen *et al.*, 1994:47; Loh *et al.*, in press:table 1.

Remarks. The presence of this species in Taiwan is questionable. It was recorded by Shen *et al.* (1993) and Chen *et al.* (1994), but the latter remarked on its similarity to *Enchelycore schismatorhynchus* and questioned whether those two species were distinct. They are in fact easily distinguished by the funnel-shaped nostril of *E. schismatorhynchus* and by the number of vertebrae, 146–153 in *E. bikiniensis* and 137–142 in *E. schismatorhynchus* (Smith, 2012).

***Enchelycore lichenosa* (Jordan & Snyder, 1901)**

苔斑勾吻鯀

*Aemasia lichenosa* Jordan & Snyder, 1901:883, fig. 20 (Type locality: Wakanoura, Wakayama Prefecture, Japan, Inland Sea, Western North Pacific). Chen & Yu, 1986:249.  
*Enchelycore lichenosa* (Jordan & Snyder, 1901): Shen *et al.*, 1993:100, fig. 15-4; Chen *et al.*, 1994:47; Loh *et al.*, in press:table 1.

Remarks. A common species found in the coral reef regions of northeastern Taiwan.

***Enchelycore pardalis* (Temminck & Schlegel, 1846)**

豹紋勾吻鯀

*Muraena pardalis* Temminck & Schlegel, 1846:268, pl. 119 (type locality: Nagasaki, Japan). Chen & Weng, 1967:15; Chen, 1969:127; Shen, 1984a:107; Chen & Yu, 1986:247; Shao *et al.*, 1994a:276.  
*Enchelycore pardalis* (Temminck & Schlegel, 1846): Shen *et al.*, 1993:101; Chen *et al.*, 1994:47; Chen, 2004:27; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

Remarks. A common species found in the coral reef regions around Taiwan.

***Enchelycore schismatorhynchus* (Bleeker, 1853)**

裂吻勾吻鯀

*Muraena schismatorhynchus* Bleeker, 1853c:301 (type locality: Bengkulu, Sumatra, Indonesia, eastern Indian Ocean).  
*Enchelycore schismatorhynchus* (Bleeker, 1853c): Shen *et al.*, 1993:101, fig. 15-6, 7; Chen *et al.*, 1994:48; Chen, 2004:27; Loh *et al.*, in press:table 1.

*Gymnothorax schismatorhynchus* (Bleeker, 1853c): Chen & Yu, 1986:247.

Remarks. A common species found in the coral reef regions of southern Taiwan.

***Enchelynassa canina* (Quoy & Gaimard, 1824)**

長牙鼻瓣鯰

*Muraena canina* Quoy & Gaimard, 1824:247 (type locality: Pulau Waigeo, Papua Barat, Indonesia and Papua New Guinea).

*Enchelynassa canina* (Quoy & Gaimard, 1824): Chen & Yu, 1986:249; Loh *et al.*, 2012:210; Loh *et al.*, in press:table 1.

Remarks. A relatively deep living rare species from southeastern Taiwan represented by only a few specimens in collections.

***Gymnomuraena zebra* (Shaw, 1797)**

斑馬裸鯰

*Gymnothorax zebra* Shaw in Shaw & Nodder, 1797: no page, pl. 322 (type locality: Sumatra, Indonesia [originally American seas]).

*Echidna zebra* (Shaw, 1797): Chen & Weng, 1967:10; Chen, 1969:127; Shen, 1984a:106; Chen & Yu, 1986:247.

*Gymnomuraena zebra* (Shaw, 1797): Shen *et al.*, 1993:101; Chen *et al.*, 1994:48; Chen, 2004:27; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

Remarks. A common species found in the coral reef regions of southern Taiwan.

***Gymnothorax albimarginatus* (Temminck & Schlegel, 1846)**

白邊裸胸鯰

*Muraena albimarginata* Temminck & Schlegel, 1846:267, pl. 118 (Type locality: Japan [Nagasaki]).

*Gymnothorax albimarginatus* (Temminck & Schlegel, 1846): Loh *et al.*, in press:table 1; Loh *et al.*, 2015:30.

?*Gymnothorax hepaticus* (not of Rüppell, 1830): Chen & Yu, 1986:249; Shen *et al.*, 1993:104; Chen *et al.*, 1994:50; Chen, 2004:28; Loh *et al.*, in press:table 1.

Remarks. A relatively deep-living and rare species represented by only a few specimens in collections. Also recorded by video taken on deep SCUBA dives at Green Island, southwestern Taiwan. *Gymnothorax hepaticus* does not occur outside the Indian Ocean, and previous records of that species from Taiwan may refer to *G. albimarginatus*.

***Gymnothorax berndti* Snyder, 1904**

班第氏裸胸鯰

*Gymnothorax berndti* Snyder, 1904:518, pl. 4, fig. 8 (type locality: Hololulu market, Oahu Island, Hawaiian Islands). Chen & Weng, 1967:29; Chen, 1969:129; Shen 1984a:108; Chen & Yu, 1986:248; Shen *et al.*, 1993:102, fig. 15-9; Chen *et al.*, 1994:49; Chen, 2004:27; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

Remarks. A relatively deep-living species that is commonly collected by hook and line at depths to 300 m around Taiwan.

***Gymnothorax bueroensis* (Bleeker, 1857)**

伯恩斯裸胸鯰

*Muraena bueroensis* Bleeker, 1857b:79 (type locality: Kajeli, Buru Island, Molucca Islands, Indonesia).

*Gymnothorax bueroensis* (Bleeker, 1857): Chen & Yu, 1986:248; Shen *et al.*, 1993:102, fig. 16-2; Chen *et al.*, 1994:49; Chen, 2004:27; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

Remarks. A common small moray, usually found in coral regions of southern Taiwan.

**Gymnothorax chilosipilus Bleeker, 1864**

唇斑裸胸鯙

*Gymnothorax chilosipilus* Bleeker, 1864b:52 (type locality: East Indian Archipelago [Indonesia]). Chen, 1956:80; Chen & Weng, 1967:36; Chen, 1969:129; Shen 1984a:108; Chen & Yu, 1986:249; Shen et al., 1993:103; Chen et al., 1994:49; Chen, 2004:27; Shao et al., 2008:238; Ho et al., 2010:table 1; Loh et al., in press:table 1.  
*Muraena nubila* (not of Richardson, 1848): Günther, 1870:117.

Remarks. A small species living on coral reefs and near shore.

**Gymnothorax chlamydatus Snyder, 1908**

黑環裸胸鯙

*Gymnothorax chlamydatus* Snyder, 1908:94 (type locality: Naha market, Okinawa Island, Ryukyu Islands, Japan). Lee & Tseng, 1985:285; Shen et al., 1993:102; Chen et al., 1994:49; Chen, 2004:27; Loh et al., in press:table 1.  
*Gymnothorax meleagris* (not of Shaw, 1795): Shen et al., 1993:104.

Remarks. A relatively deep-living and rare species that is only represented by a few specimens in collections.

**Gymnothorax elegans Bliss, 1883**

雅緻裸胸鯙

*Gymnothorax elegans* Bliss, 1883:60 (type locality: Mauritius, Mascarenes, southwestern Indian Ocean). Loh et al., 2012:211.

Remarks. A rare species living in deep water and represented by a few specimens in collections in Taiwan.

**Gymnothorax eurostus (Abbott, 1860)**

徽身裸胸鯙

*Thyrsoidea eurosta* Abbott, 1860:478 (type locality: Hawaiian Islands).

*Gymnothorax eurostus* (Abbott, 1860): Shen et al., 1993:103; Chen et al., 1994:50; Chen et al., 1997:appendix 1; Shao et al., 2008:238; Ho et al., 2010:table 1; Loh et al., in press:table 1.

*Gymnothorax meleagris* (not of Shaw, 1795): Chen & Weng, 1967:20; Shen et al., 1993:103.

Remarks. A common species is found in coral reef regions around Taiwan. It is similar to *G. meleagris*, and the two species have been confused in the past (i.e. Chen & Weng, 1967). In a further complication, Hatooka (2002:1453) concluded that the holotype of *G. meleagris* is actually the species currently known as *G. eurostus*. Smith (2012) examined this specimen and found the identification difficult to resolve. Further studies are needed, but we retain the traditional interpretation here.

**Gymnothorax favagineus Bloch & Schneider, 1801**

黑斑裸胸鯙

*Gymnothorax favagineus* Bloch & Schneider, 1801:525, pl. 105 (type locality: Tharangambadi, India). Chen, 1956:80; Chen & Weng, 1967:33; Chen, 1969:129; Chen & Yu, 1986:249; Chen et al., 1994:50; Shao et al., 1994b:64; Shen et al., 1993:103; Chen, 2004:28; Chen & Loh 2007:80; Shao et al., 2008:238; Ho et al., 2010:table 1; Loh et al., in press:table 1.

Remarks. This species is easily confused with *G. isingteena*, and the records above might include misidentification of that species. The color pattern seems to change with growth and further study is needed to confirm its identification.

**Gymnothorax fimbriatus (Bennett, 1832)**

縫斑裸胸鯙

*Muraena fimbriata* Bennett, 1832:168 (type locality: Mauritius, Mascarenes, southwestern Indian Ocean).

*Gymnothorax fimbriatus* (Bennett, 1832): Jordan & Richardson, 1909:173; Chen & Weng, 1967:32; Chen, 1969:129; Chen & Yu, 1986:249; Shen, 1988:12; Shen et al., 1993:103; Chen et al., 1994:50; Chen et al., 1997:appendix 1; Chen, 2004:28; Shao et al., 2008:238; Ho et al., 2010:table 1; Loh et al., in press:table 1.

Remarks. A common species found in the coral reef regions around Taiwan.

***Gymnothorax flavimarginatus* (Rüppell, 1830)**

黃邊鰭裸胸鯙

*Muraena flavimarginata* Rüppell, 1830:119, pl. 30, fig. 3 (type locality: Red Sea).

*Gymnothorax flavimarginatus* (Rüppell, 1830): Jordan & Richardson, 1909:240; Chen, 1956:80; Chen & Weng, 1967:22; Chen, 1969:127; Shen, 1984a:107; Chen & Yu, 1986:248; Shen et al., 1993:103; Chen et al., 1994:50; Chen, 2004:28; Shao et al., 2008:238; Ho et al., 2010:table 1; Loh et al., in press:table 1.

Remarks. A species commonly found in the coral reef regions around Taiwan.

***Gymnothorax formosus* Bleeker, 1864**

美麗裸胸鯙

*Gymnothorax formosus* Bleeker, 1864b:51 (type locality: Wahai, Ceram, Indonesia). Loh et al., in press:table 1.

Remarks. A species commonly found in the coral reef regions of southeastern Taiwan.

***Gymnothorax fuscomaculatus* (Schultz, 1953)**

褐點裸胸鯙

*Rabula fuscomaculata* Schultz in Schultz, et al., 1953:147, fig. 30 (Type locality: Marshall Islands, Western Pacific).

*Gymnothorax fuscomaculatus* (Schultz, 1953): Loh et al., 2012:213.

Remarks. A very rare species represented by only one specimen collected from southern Taiwan.

***Gymnothorax gracilicauda* Jenkins, 1903**

細尾裸胸鯙

*Gymnothorax gracilicauda* Jenkins, 1903:426, fig. 6 (type locality: Honolulu, Oahu Island, Hawaiian Islands). Loh et al., in press:table 1.

Remarks. A very rare species in Taiwan.

***Gymnothorax herrei* Beebe & Tee Van, 1933**

海氏裸胸鯙

*Gymnothorax brunneus* Herre, 1923:212 (type locality: Analao, Batangas, Philippines, 13°45'N, 120°55'E, 0–3 m) (Preoccupied). Shen et al., 1993:102, fig. 15-10; Chen et al., 1994:49.

*Gymnothorax herrei* Beebe & Tee Van, 1933:138 (replacement name for *Gymnothorax brunneus* Herre, 1923). Shao et al., 2008:238; Loh et al., in press:table 1.

Remarks. There is only one specimen collected from a tide pool in southern Taiwan.

***Gymnothorax intesi* (Fourmanoir & Rivaton, 1979)**

英氏裸胸鯙

*Lycodontis intesi* (Fourmanoir & Rivaton, 1979):426, fig. 16 (type locality: Lifou Island, Loyalty Islands, southwestern Pacific, 400 m).

*Gymnothorax intesi* Fourmanoir & Rivaton, 1979: Loh et al., in press:table 1.

Remarks. A deep water species that occasionally collected by bottom trawl or hook and line off southeastern Taiwan.

### ***Gymnothorax isingteena* (Richardson, 1845)**

大斑裸胸鯙

*Muraena isingteena* Richardson, 1845a: pl. 48, fig. 1 (type locality: Canton, China).

*Gymnothorax isingteena*: (Richardson, 1845): Ho & Shao, 2011:21, table 1; Smith 2012:17.

*Gymnothorax melanospilos* (Bleeker, 1855): Jordan & Richardson, 1909:174, fig. 67; Okada, 1938:155; Chen, 1956:80; Chen & Weng, 1967:30; Chen, 1969:129; Shen, 1984b:109; Chen & Yu, 1986:248; Shen, 1988:13.

\**Gymnothorax pescadoris* Jordan & Evermann, 1902:173, fig. 8 (type locality: Kotosho, Hokoto [Taipai City] or Pescadores Islands, Taiwan). Jordan & Richardson, 1909:326, fig. 9; Okada & Matsubara, 1938:99, pl.14, fig. 3; Matsubara, 1955:354; Chen & Weng, 1967:31; Chen, 1969:129; Chen & Yu, 1986:249; Shao et al., 1994a:276.

Remarks. This species is easily confused with *G. favagineus*, and the citations provided above might include misidentifications of that species. Also see Remarks under *G. favagineus*. Common in Taiwan.

### ***Gymnothorax javanicus* (Bleeker, 1859)**

爪哇裸胸鯙

*Muraena javanica* Bleeker, 1859:347 (type locality: Patjitan, Java, Indonesia).

*Gymnothorax javanicus* (Bleeker, 1959): Shen et al., 1993:104; Chen et al., 1994:50; Chen et al., 1997:appendix 1; Chen, 2004:28; Shao et al., 2008:238; Loh et al., in press:table 1.

Remarks. A common species collected by hook and line from coral reef areas around Taiwan. It is the largest of all moray eels by weight.

### ***Gymnothorax kidako* (Temminck & Schlegel, 1846)**

蟠紋裸胸鯙

*Muraena kidako* Temminck & Schlegel, 1846:266, pl. 117 (type locality: Nagasaki, Japan).

*Gymnothorax kidako* (Temminck & Schlegel, 1846): Chen, 1956:80; Chen & Weng, 1967:25; Chen, 1969:128; Shen 1984a:108; Chen & Yu, 1986:248; Shen et al., 1993:104; Chen et al., 1994:51; Chen, 2004:28; Shao et al., 2008:238; Loh et al., in press:table 1.

Remarks. A common species collected by hook and line at shallow waters around Taiwan.

### ***Gymnothorax margaritophorus* Bleeker, 1864**

斑頸裸胸鯙

*Gymnothorax margaritophorus* Bleeker, 1864b:53 (type locality: Ambon Island, Molucca Islands, Indonesia). Shen et al., 1993:104; Chen & Wang, 1994a:159; Chen et al., 1994:51; Chen, 2004:28; Shao et al., 2008:238; Ho et al., 2010:table 1; Loh et al., in press:table 1.

Remarks. A small species common on coral reefs.

### **\**Gymnothorax melanosomatus* Loh, Shao & Chen, 2011**

黑身裸胸鯙

*Gymnothorax melanosomatus* Loh, Shao & Chen, 2011:44, figs. 1–2 (type locality: offshore from Changbin, Taitung, Taiwan, 50–100 m). Smith, 2012:19; Loh et al., in press:table 1; Loh et al., 2015:30.

Remarks. A relatively deep-living species that is commonly collected by hook and line off southeastern Taiwan.

**Gymnothorax melatremus Schultz, 1953**

黃身裸胸鯙

*Gymnothorax melatremus* Schultz in Schultz, et al., 1953:120, figs. 23d, 25 (type locality: Boby Island, Bikini Atoll, Marshall Islands, western Pacific). Shen et al., 1993:104; Chen et al., 1994:51; Loh et al., in press:table 1.

Remarks. A small species with small body size collected around coral reefs.

**Gymnothorax meleagris (Shaw, 1795)**

白口裸胸鯙

*Muraena meleagris* Shaw in Shaw & Nodder, 1795: no page number, pl. 220 (type locality: Southern Ocean).

*Gymnothorax meleagris* (Shaw, 1795): Chen, 1956:79; Chen & Weng, 1967:20; Shen, 1984a:107; Shen, 1984:106; Chen & Yu, 1986:247; Shen et al., 1993:105; Chen et al., 1994:51; Chen et al., 1997:appendix 1; Chen, 2004:29; Shao et al., 2008:238; Loh et al., in press:table 1.

*Gymnothorax chlorostigma* (Kaup, 1856): Chen, 2004:27.

Remarks. See Remarks under *G. eurostus*.

**Gymnothorax minor (Temminck & Schlegel, 1846)**

小裸胸鯙

*Muraena minor* Temminck & Schlegel, 1846:269, pl. 115, fig. 2 (type locality: Japan).

*Gymnothorax minor* (Temminck & Schlegel, 1846): Chen, 2004:29; Shao et al., 2008:238; Ho et al., 2010:table 1; Loh et al., in press:table 1.

*Gymnothorax reticularis* (not of Bloch, 1795): Jordan & Richardson, 1909:173; Chen, 1956:80; Chu, 1957:15; Liang & Yeh, 1964:8; Lee, 1967:44; Chen & Weng, 1967:23; Chen, 1969:127; Shen, 1984a:107; Chen & Yu, 1986:248; Shen, 1988:14; Shen et al., 1993:106; Chen et al., 1994:52; Shao et al., 1994a:276; Shao et al., 2008:238.

*Lycodontis reticularis* (not of Bloch, 1795): Fowler, 1932b:137, fig. 12.

Remarks. The most abundant moray species collected by bottom trawl off Taiwan. Previously referred to *G. reticularis* by many authors.

**Gymnothorax monostigma (Regan, 1909)**

眼斑裸胸鯙

*Muraena monostigma* Regan, 1909:438 (type locality: Raiatea, Society Islands).

*Gymnothorax monostigma* (Regan, 1909): Chen & Yu, 1986:249; Shao et al., 2008:238; Loh et al., in press:table 1.

*Gymnothorax monostigmus* (Regan, 1909): Shen et al., 1993:105.

Remarks. A rare species collected by hook and line off southeastern Taiwan.

**\*Gymnothorax neglectus Tanaka, 1911**

細花斑裸胸鯙

*Gymnothorax neglectus* Tanaka, 1911:28, pl. 7, fig. 24 (type locality: Keelung, Taihoku, northern Taiwan). Chen & Weng, 1967:35; Chen, 1969:129; Chen & Yu, 1986:249; Shen et al., 1993:105; Chen et al., 1994:51; Shao et al., 2008:238; Ho et al., 2010:table 1; Ho & Shao, 2011:21, table 1; Smith, 2012:21; Loh et al., in press:table 1.

Remarks. A relatively deep living species that is commonly collected by hook and line down to 300 m depth, in eastern Taiwan.

**\**Gymnothorax niphostigmus* Chen, Shao & Chen, 1996**

雪花斑裸胸鯥

*Gymnothorax niphostigmus* Chen, Shao & Chen, 1996:20, figs. 1–3 (type locality: Hopingtao fish market, Keelung, Taiwan, 100–150 m). Chen, 2004:29; Ho & Shao, 2011:21, table 1; Smith, 2012:22; Loh *et al.*, in press:table 1.

Remarks. A relatively deep-living species commonly collected by hook and line off eastern Taiwan.

***Gymnothorax nudivomer* (Günther, 1867)**

裸鋤裸胸鯥

*Muraena nudivomer* Günther in Playfair & Günther, 1867:127, pl. 18 (type locality: Zanzibar, Tanzania, western Indian Ocean).

*Gymnothorax nudivomer* (Günther, 1867): Shen *et al.*, 1993:105; Chen *et al.*, 1994:52; Chen, 2004:29; Loh *et al.*, in press:table 1.

Remarks. A common species found in coral reef regions around Taiwan.

***Gymnothorax phasmatodes* (Smith, 1962)**

無斑裸胸鯥

*Lycodontis phasmatodes* Smith, 1962:436, pl. 53, figs B, C & I (type locality: Inhaca Island, Mozambique, western Indian Ocean).

*Gymnothorax phasmatodes* (Smith, 1962): Loh *et al.*, in press:table 1; Loh *et al.*, 2015:30.

Remarks. A small species that is commonly collected by hook and line off southern Taiwan.

***Gymnothorax pictus* (Ahl, 1789)**

細點裸胸鯥

*Muraena picta* Ahl, 1789:8, pl. 2, right fig. (type locality: in Indian Orientalis [Indonesia]).

*Gymnothorax pictus* (Ahl, 1789): Chen & Weng, 1967:19; Chen, 1969:127; Shen, 1984a:107; Chen & Yu, 1986:247; Shao *et al.*, 1994a:276; Chen *et al.*, 1994:53; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

*Siderea picta* (Ahl, 1789): Chen, 2004:29; Shen *et al.*, 1993:107.

Remarks. A common species found in coral reef regions around Taiwan.

***Gymnothorax pindae* Smith, 1962**

平達裸胸鯥

*Gymnothorax pindae* Smith, 1962:430, pl. 55, fig. D (type locality: Pinda, Mozambique, western Indian Ocean). Shen *et al.*, 1993:105; Chen *et al.*, 1994:52; Shao *et al.*, 2008:238; Loh *et al.*, in press:table 1.

Remarks. A rare species only found in coral reef regions of southern Taiwan.

***Gymnothorax polyuranodon* (Bleeker, 1854)**

豹紋裸胸鯥

*Muraena polyuranodon* Bleeker, 1854:248 (type locality: Ceram, Indonesia).

*Gymnothorax polyuranodon* (Bleeker, 1854): Chen, 1956:79; Chen & Weng, 1967:21; Chen, 1969:127; Shen, 1984a:107; Chen & Yu, 1986:248; Shao *et al.*, 2008:238; Loh *et al.*, in press:table 1.

Remarks. A rare species found in freshwater regions around northeastern and southwestern Taiwan.

***Gymnothorax prionodon* Ogilby, 1895**

鋸齒裸胸鯡

*Gymnothorax prionodon* Ogilby, 1895:720 (type locality: Port Jackson, New South Wales, Australia). Ho *et al.*, 2010:table 1; Ho & Shao, 2011:21, table 1; Smith, 2012:25.

\**Gymnothorax leucostigma* Jordan & Richardson, 1909:174, pl. 68 (type locality: Kaohsiung, Taiwan). Okada, 1938:154; Chen, 1956:80; Chen & Weng, 1967:34; Chen, 1969:129; Chen & Yu, 1986:249; Chen *et al.*, 1994:51; Shen *et al.*, 1993:104;

*Gymnothorax leucostigmus* (Jordan & Richardson, 1909): Chen, 2004:28.

Remarks. A relatively deep living species that is commonly collected by hook and line off southeastern Taiwan.

**\**Gymnothorax prolatus* Sasaki & Amaoka, 1991**

小長鯡

*Gymnothorax prolatus* Sasaki & Amaoka, 1991:7, figs. 1–3 (type locality: Suao fish market, northeastern Taiwan). Shao *et al.*,

2008:238; Ho *et al.*, 2010:table 1; Ho & Shao, 2011:22, table 1; Loh *et al.* 2011:45; Smith, 2012:25; Loh *et al.*, 2015:30.

*Strophidion prolatus* (Sasaki & Amaoka, 1991): Loh *et al.*, in press:table 1.

Remarks. A deep living species that is occasionally collected by bottom trawls off northeastern and southwestern Taiwan.

**\**Gymnothorax pseudomelanosomatus* Loh, Shao & Chen, 2015**

擬黑身裸胸鯡

*Gymnothorax pseudomelanosomatus* Loh, Shao & Chen in Loh *et al.*, 2015:32. (type locality: Changbin, Taitung county, southeastern Taiwan).

Remarks. A deep-living species that is only represented by the two type specimens collected from southeastern Taiwan and one specimen from southwestern Taiwan (USNM 395050); it is easily confused with *Gymnothorax melanosomatus*, which was also described from same region.

***Gymnothorax pseudothyrsoides* (Bleeker, 1853)**

淡網紋裸胸鯡

*Muraena pseudothyrsoides* Bleeker, 1853b:778 (type locality: Makassar, Sulawesi, Indonesia).

*Gymnothorax pseudothyrsoides* (Bleeker, 1853b): Jordan & Richardson, 1909:173; Chen, 1956:80; Chen & Weng, 1967:28; Chen, 1969:129; Shen 1984a:108; Shen, 1984b:12; Chen & Yu, 1986:248; Shen *et al.*, 1993:105; Chen *et al.*, 1994:52; Chen *et al.*, 1997:appendix 1; Shao *et al.*, 1994a:276; Chen, 2004:29; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

Remarks. A common species found in coral reef regions around Taiwan.

***Gymnothorax reevesii* (Richardson, 1845)**

雷福氏裸胸鯡

*Muraena reevesii* Richardson, 1845a: pl. 49, fig. 2 (type locality: China seas).

*Gymnothorax reevesii* (Richardson, 1845): Shen *et al.*, 1993:106; Chen *et al.*, 1994:52; Chen, 2004:29; Loh *et al.*, in press:table 1.

Remarks. A common species found in coral reef regions around Taiwan.

***Gymnothorax richardsonii* (Bleeker, 1852)**

李氏裸胸鯡

*Muraena richardsonii* Bleeker, 1852:296 (type locality: Wahai, northern Ceram and Padang, western Sumatra, Indonesia).

*Gymnothorax richardsonii* (Bleeker, 1852): Shao *et al.*, 2008:238; Loh *et al.*, in press:table 1.

*Gymnothorax richardsoni* (Bleeker, 1852): Chen & Yu, 1986:248; Shen et al., 1993:106; Chen et al., 1994:52.

Remarks. An uncommon species found in southeastern and southern Taiwan occasionally.

***Gymnothorax rueppelliae* (McClelland, 1844)**

寬帶裸胸鯙

*Dalophis rueppelliae* McClelland, 1844:213 (type locality: Red Sea).

*Gymnothorax rueppelliae* (McClelland, 1844): Shen et al., 1993:106; Chen et al., 1994:53; Chen, 2004:29; Loh et al., in press:table 1.

*Gymnothorax petelli* (Bleeker, 1856): Chen & Yu, 1986:248.

*Gymnothorax ruppelli* (McClelland, 1844): Chen & Yu, 1986:248.

*Gymnothorax ruppelli* (McClelland, 1844): Shao et al., 2008:238.

Remarks. A common species found in coral reef regions around Taiwan.

***Gymnothorax sagmacephalus* Böhlke, 1997**

鞍斑裸胸鯙

*Gymnothorax sagmacephalus* Böhlke, 1997:100, figs. 12–14 (type locality: Tokyo Bay, Japan). Loh et al., in press:table 1; Loh et al., 2015:30.

Remarks. A small to moderate-size species that is commonly collected by hook and line off southern Taiwan.

**\**Gymnothorax shaoi* Chen & Loh, 2007**

邵氏裸胸鯙

*Gymnothorax shaoi* Chen & Loh, 2007:77, figs. 1–4 (type locality: off shore of Sanshengtai, 23.130°N, 121.414°E, Chengkung, Taitung, Taiwan.). Ho & Shao, 2011:22, table 1; Smith, 2012:27; Loh et al., in press:table 1.

Remarks. A species commonly collected by hook and line in coral reef regions of southeastern Taiwan.

**\**Gymnothorax taiwanensis* Chen, Loh & Shao, 2008**

臺灣裸胸鯙

*Gymnothorax taiwanensis* Chen, Loh & Shao, 2008:132, figs. 1–2 (type locality: offshore from Shihtiping, 23°29'N, 121°30'E, Hualien, eastern Taiwan, 10 m). Ho & Shao, 2011:22, table 1; Smith, 2012:28; Loh et al., in press:table 1.

Remarks. A species commonly collected by hook and line in the coral reef regions of southeastern Taiwan.

***Gymnothorax thyrsoideus* (Richardson, 1845)**

密點裸胸鯙

*Muraena thrysoidea* Richardson, 1845b:111 (type locality: Canton, China).

*Gymnothorax thyrsoideus* (Richardson, 1845): Shao et al., 2008:238; Ho et al., 2010:table 1; Loh et al., in press:table 1.

*Gymnothorax thyrsoidea* (Richardson, 1845): Chen & Yu, 1986:247; Shen et al., 1993:107.

*Siderea thyrsoidea* (Richardson, 1845): Chen et al., 1994:52; Chen, 2004:30.

Remarks. A common species found in coral reef regions around Taiwan.

***Gymnothorax undulatus* (Lacepède, 1803)**

疏斑裸胸鯙

*Muraenophis undulata* Lacepède (ex. Commerson), 1803:629, 642, pl. 19, fig. 2 (no type locality stated).

*Gymnothorax undulatus* (Lacepède, 1803): Chen, 1956:80; Chen & Weng, 1967:27; Chen, 1969:129; Shen 1984a:108; Chen &

Yu, 1986:248; 1991:140; Shen *et al.*, 1993:106; Chen *et al.*, 1994:53; Chen, 2004:29; Shao *et al.*, 2008:238; Loh *et al.*, in press:table 1.

Remarks. A common species found in coral reef regions around Taiwan.

***Gymnothorax ypsilon* Hatooka & Randall, 1992**

丫環裸胸鯀

*Gymnothorax ypsilon* Hatooka & Randall, 1992:183, figs. 1–2 (type locality: Uotsurijima Island, Senkaku Islands, Japan). Chen, 2004:29; Loh *et al.*, in press:table 1.

*Gymnothorax punctatofasciatus* (not of Bleeker, 1863): Okada, 1938:154; Chen, 1956:80; Chen & Weng, 1967:24; Chen, 1969:127; Shen, 1984a:108; Chen & Yu, 1986:248.

Remarks. This rare species is found in relatively deep coral reef regions around Taiwan. It was confused with *G. punctatofasciatus* by previous authors. The records of *G. punctatofasciatus* in Taiwan refers to this species.

***Gymnothorax zonipectis* Seale, 1906**

帶尾裸胸鯀

*Gymnothorax zonipectis* Seale, 1906:7, fig. 1 (type locality: Tahiti, Society Islands, South Pacific). Shen *et al.*, 1993:106; Chen *et al.*, 1994:53; Chen *et al.*, 1997:appendix 1; Shao *et al.*, 2008:238; Loh *et al.*, in press:table 1.

Remarks. A rare species found in coral reef regions of northeastern and southern Taiwan.

***Pseudechidna brummeri* (Bleeker, 1858)**

布氏擬蝮鯀

*Muraena brummeri* Bleeker, 1858a:137 (type locality: Atapupu, Timor Island, southern Malay Archipelago).

*Pseudechidna brummeri* (Bleeker, 1858): Loh *et al.*, in press:table 1; Loh *et al.*, 2015:30.

*Strophidion brummeri* (Bleeker, 1858): Shen 1984a:110; Chen & Yu, 1986:249; Shen *et al.*, 1993:107; Chen *et al.*, 1994:54.

Remarks. A rare species found in coral reef regions of southeastern Taiwan.

***Rhinomuraena quaesita* Garman, 1888**

管鼻鯀

*Rhinomuraena quaesita* Garman, 1888:114, fig. (type locality: Ebon Island, Marshall Islands, western Pacific). Shen, 1974:181; Shen, 1984a:106; Chen & Yu, 1986:247; Shen *et al.*, 1993:107; Chen *et al.*, 1994:53; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

Remarks. A rare species found in coral reef regions of southeastern Taiwan.

***Scuticaria tigrina* (Lesson, 1828)**

虎斑長圓鯀

*Ichthyophis tigrinus* Lesson, 1828:399 (type locality: Bora Bora, Society Islands [French Polynesia]).

*Uropterygius tigrina* (Lesson, 1828): Shen *et al.*, 1993:99.

*Scuticaria tigrina* (Lesson, 1828): Chen, 2004:29; Shao *et al.*, 2008:238; Loh *et al.*, in press:table 1.

Remarks. A rare species found in coral reef regions of southeastern Taiwan.

***Strophidion dorsalis* (Seale, 1917)**

長背長鯀

*Gymnothorax dorsalis* Seale, 1917:92 (type locality: Hong Kong). Loh *et al.*, in press:table 1.

*Strophidon dorsalis* (Seale, 1917): Loh *et al.*, 2015:30.

Remarks. This deep water species is commonly collected by bottom trawl or hook and line around Taiwan.

### ***Strophidon sathete* (Hamilton, 1822)**

長鰐

*Muraenophis sathete* Hamilton, 1822:17, 363 (type locality: Ganges estuaries near Calcutta, India).

*Strophidon sathete* (Hamilton, 1822): Shen *et al.*, 1993:108; Chen *et al.*, 1994:54; Shao *et al.*, 2008:238; Loh *et al.*, in press:table 1; Loh *et al.*, 2015:30.

*Thyrsoidea macrurus* (Bleeker, 1854): Chen, 1956:79; Chen & Weng, 1967:14; Chen, 1969:127; Chen & Yu, 1986:247.

*Strophidon ui* (Tanaka, 1918): Chen & Weng, 1967:37; Chen, 1969:129; Chen & Yu, 1986:249.

*Thyrsoidea macrurus* (Bleeker, 1854): Shen, 1984a:106.

*Evenchelys macrurus* (Bleeker, 1854): Jordan & Richardson, 1902:327; Fowler, 1932b:144, fig. 16.

Remarks. A very common species collected by bottom trawl and traps. This is the largest moray in length, though not in weight, with a total length up to 3.7 meters (Randall *et al.*, 1997).

### **+*Uropterygius alboguttatus* Smith, 1962**

白斑鰭尾鯽

*Uropterygius alboguttatus* Smith, 1962:427, pl. 60, fig. E (type locality: Assumption Island, western Indian Ocean).

*Uropterygius xanthopterus* (not of Bleeker, 1859): Ho *et al.*, 2010:table 1.

Voucher specimen: NMNM-P 2560 (Taiwan).

Remarks. The specimen of *Uropterygius xanthopterus* listed in Ho *et al* (2010) is reidentified as this species. This is a new record in Taiwan.

### ***Uropterygius macrocephalus* (Bleeker, 1864)**

巨頭鰭尾鯽

*Gymnomuraena macrocephalus* Bleeker, 1864b:54 (type locality: Ambon Island, Molucca Islands, Indonesia).

*Uropterygius macrocephalus* (Bleeker, 1864b): Shen *et al.*, 1993:99; Chen *et al.*, 1994:62; Chen *et al.*, 1997:appendix 1; Loh *et al.*, 2008:140; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1; Loh *et al.*, in press:table 1.

Remarks. A small species found in coral reef regions around Taiwan.

### ***Uropterygius marmoratus* (Lacepède, 1803)**

石點長圓鯽

*Gymnomuraena marmorata* Lacepède (ex Commerson), 1803:648, 649 (type locality: New Britain Island, Bismarck Archipelago, Papua New Guinea, western Pacific). Chen & Yu, 1986:249.

*Scuticaria marmorata* (Lacepède, 1803): Loh *et al.*, 2008:140; Loh *et al.*, in press:table 1.

Remarks. A rare species found in coral reef regions of southeastern Taiwan. Loh *et al.* (2008) transferred this species from its traditional placement in the genus *Uropterygius* to *Scuticaria*. This placement requires further research.

### ***Uropterygius micropterus* (Bleeker, 1852)**

小鰭鰭尾鯽

*Muraena micropterus* Bleeker, 1852:298 (type locality: Wahai, northern Ceram, Indonesia).

*Uropterygius micropterus* (Bleeker, 1852): Shen *et al.*, 1993:99; Chen *et al.*, 1994:62; Chen, 2004:30; Shao *et al.*, 2008:238; Loh *et al.*, 2008:140; Loh *et al.*, in press:table 1.

Remarks. A small species found in coral reef regions around Taiwan.

***Uropterygius nagoensis* Hatooka, 1984**

網紋鰭尾鯉

*Uropterygius nagoensis* Hatooka, 1984:20, figs. 1–4 (type locality: Nago fish market, Okinawa Island, Ryukyu Islands, western North Pacific). Chen, 2004:30.

Remarks. A very widely distributed species in the Indo-west Pacific region, but only recorded in Penghu Island in Taiwan.

**\**Uropterygius oligospondylus* Chen, Randall & Loh, 2008**

寡椎鰭尾鯉

*Uropterygius oligospondylus* Chen, Randall & Loh in Loh *et al.*, 2008:141, figs. 1H, 3–4 (type locality: Changbin, Taitung, southeastern Taiwan, 15 m). Ho & Shao, 2011:22, table 1; Smith, 2012:37; Loh *et al.*, in press:table 1.

Remarks. A medium-size species found in coral reef regions of southeastern Taiwan.

***Uropterygius xanthopterus* Bleeker, 1859**

黃鰭鰭尾鯉

*Uropterygius xanthopterus* Bleeker, 1859:350 (type locality: Patjitan, Java, Indonesia).

Voucher specimens: USNM 263107, 312868 (Kenting, S. Taiwan).

Remarks. The only specimen listed in Ho *et al.* (2010) was subsequently reidentified as *U. alboguttatus* by DGS. There are two other specimens in the USNM collection, however, that confirm the presence of the species.

**Family Nemichthyidae**

線鰻科

***Avocettina infans* (Günther, 1878)**

喙吻鰻

*Nemichthys infans* Günther, 1878:251 (type locality: Mid-Atlantic, 5°48'N, 14°20'W, 2500 fathoms [4572 m]).

*Avocettina infans* (Günther, 1878): Chen & Yu, 1986:261 (based on Nielsen & Smith, 1978). Shao *et al.*, 2008:239.

Remarks. A common species collected by midwater trawl or bottom trawl off northeastern and southwestern Taiwan. It is easily confused with *Nemichthys scolopaceus* but can be distinguished by the dark body color, the anus well behind the pectoral fin, and the non-filamentous tail.

***Nemichthys scolopaceus* Richardson, 1848**

線鰻

*Nemichthys scolopaceus* Richardson, 1848a:25, pl. 10, figs. 1–3 (type locality: south Atlantic off Brazil). Chen & Weng, 1967:84; Shen, 1984a:106; Chen & Yu, 1986:261; Shen *et al.*, 1993:108; Shao *et al.*, 2008:239; Ho *et al.*, 2010:table 1.

Remarks. A very common species collected by midwater or bottom trawls off northeastern and southwestern Taiwan.

## **Family Nettastomatidae**

鴨嘴鰻科

### ***Nettastoma parviceps* Günther, 1877**

小頭鴨嘴鰻

*Nettastoma parviceps* Günther, 1877:446 (type locality: south of Tokyo, Japan, 345 fathoms [631 m]). Shao *et al.*, 2008:239.

Remarks. An uncommon deepwater species collected by bottom trawl off northeastern Taiwan.

### ***Nettastoma solitarium* Castle & Smith, 1981**

前鼻鴨嘴鰻

*Nettastoma solitarium* Castle & Smith in Smith, Böhlke & Castle, 1981:548, figs. 2–7 (Cabra Island, the Philippines, western Pacific, 415–510 m). Shao *et al.*, 2008:239; Ho *et al.*, 2010:table 1.

Remarks. A deepwater species occasionally collected by bottom trawl around Taiwan.

### **+*Nettenchelys gephyra* Castle & Smith, 1981**

橋鴨蛇鰻

*Nettenchelys gephyra* Castle & Smith in Smith, Böhlke & Castle, 1981:553, figs. 8–11 (type locality: New Caledonia). Smith *et al.*, 2015:110.

Remarks. A rare deep-sea species collected by bottom trawl off northeastern Taiwan. This is a new record in Taiwan.

### **\**Nettenchelys proxima* Smith, Lin & Chen, 2015**

近鴨蛇鰻

*Nettenchelys proxima* Smith, Lin & Chen in Smith *et al.*, 2015:114. (type locality: Dong-gang, southwestern Taiwan).

Remarks. A rare species only represented by the holotype and a tentatively identified specimen from the Philippines.

### ***Saurenchelys fierasfer* (Jordan & Snyder, 1901)**

線尾鴨嘴鰻

*Chlopsis fierasfer* Jordan & Snyder, 1901:860, fig. 10 (type locality: Wakanoura, Wakayama Prefecture, Japan). Chen, 1956:77; Liaw, 1960:4; Lee, 1967:44; Chen & Weng, 1967:80; Chen, 1969:134; Shen 1984a:114; Chen & Yu, 1986:255; Shen *et al.*, 1993:115.

*Saurenchelys fierasfer* (Jordan & Snyder, 1901): Shen, 1988:15; Shao *et al.*, 1994a:276; Chen, 2004:32; Shao *et al.*, 2008:239; Ho *et al.*, 2010:table 1.

Remarks. A very common species collected by midwater or bottom trawls around Taiwan.

### **\**Saurenchelys gigas* Lin, Smith & Shao, 2015**

巨蜥鰻

*Saurenchelys gigas* Lin, Smith & Shao In Lin *et al.*, 2015:122 (type locality: Daxi, Yilan, northeastern Taiwan).

Remarks. This is the largest known species of the genus and is known from Taiwan and Vietnam.

### **+*Saurenchelys taiwanensis* Karmovskaya, 2004**

台灣蜥鰻

*Saurenlchelys taiwanensis* Karmovskaya, 2004:S27 (Type locality: Philippines).

Voucher specimens: NMMB-P 10064, 10065, 11190, 11193, 11986, 12624, 13160, 15516, 15911, 16451, 19446 (Dong-gang); NMMB-P 12229 (Da-xi).

Remarks. This species was originally described from a single specimen collected in the Philippines. There are many specimens newly collected from deep waters off northeastern and southwestern Taiwan. This is a new record in Taiwan.

## Family Ophichthidae

蛇鰐科

### *Bascanichthys kirkii* (Günther, 1870)

褐蛇鰐

*Ophichthys kirkii* Günther, 1870:89 (type locality: Rovuma Bay, eastern Africa).

*Bascanichthys kirkii* (Günther, 1870): Chen & Weng, 1967:62; Chen, 1969:137; Shen 1984a:116; Chen & Yu, 1986:258; Shen et al., 1993:110; Shao et al., 2008:238; Ho et al., 2010:table 1.

Remarks. An uncommon species; several specimens have been collected from river mouths around Taiwan.

### *Brachysomophis cirrocheilos* (Bleeker, 1857)

大口短體鰐

*Ophisurus cirrocheilos* Bleeker, 1857a:89 (type locality: Ambon, Indonesia).

*Ophichthus cirrocheilos* (Bleeker, 1857a): Günther, 1870:65.

*Brachysomophis cirrocheilos* (Bleeker, 1857a): Chen & Weng, 1967:58; Chen, 1956:75; Chen, 1969:135; Shen 1984a:116; Chen & Yu, 1986:258; Shen et al., 1993:111; McCosker & Randall, 2001:8; Shao et al., 2008:238; McCosker & Ho, 2014:95.

Remarks. A common species collected from muddy bottoms in shallow waters of western and southwestern Taiwan. It is sometimes confused with *Ophichthus lithinus* because of their similar coloration.

### +*Brachysomophis crocodilinus* (Bennett, 1833)

鰐形短體鰐

*Ophisurus crocodilinus* Bennett, 1833:32 (type locality: Ambon Island, Molucca Islands, Indonesia).

Voucher specimens: Two specimens in the TOU-AE collection.

Remarks. This is a new record for Taiwan. It is commonly observed over sandy bottom just outside the coral reef region. Underwater photographs of the species were examined and their identification confirmed by HH. A few specimens were recently collected from southeastern Taiwan.

### +*Brachysomophis henshawi* Jordan & Snyder, 1904

亨氏短體鰐

*Brachysomophis henshawi* Jordan & Snyder, 1904:940 (type locality: Honolulu, Hawaiian Is.).

Voucher specimens: FRIP 21492, 22153 (Taitung, southeastern Taiwan).

Remarks. This is a new record for Taiwan. Known only from several specimens collected by hook and line off southeastern Taiwan.

### \**Brachysomophis longipinnis* McCosker & Randall, 2001

長鰭短體蛇鰐

*Brachysomophis longipinnis* McCosker & Randall, 2001:23, figs. 5, 7–8, 13 (type locality: Taiwan Strait, ca. 50 m). Ho & Shao, 2011:23, table 1; McCosker & Ho, 2014:96.

Remarks. This species was originally described from the single holotype. McCosker & Ho (2014) reported two additional specimens recently collected in southwestern Taiwan.

**+*Brachysomophis porphyreus* ((Temminck & Schlegel, 1846)**

紫身短體蛇鰻

*Ophisurus porphyreus* Temminck & Schlegel, 1846:265, pl. 116 (type locality: Japan).

*Brachysomophis porphyreus* (Temminck & Schlegel, 1846): McCosker & Randall 2001:25.

Voucher specimens: ASIZP 70365, NMMB-P10465 (Da-xi); ASIZP 72251 (Dong-shi).

Remarks. Although McCosker & Randall (2001) noted that this species occurs in the Taiwan Strait, no voucher was available. Recently, several specimens were collected by the first author from shallow water near a river mouth in the Taiwan Strait. It attains a large size, with total length approaching 2 m.

***Callechelys kuro* (Kuroda, 1947)**

黑麗鰻

*Aphthalmichthys kuro* Kuroda, 1947:27 (type locality: Tosa Bay, Japan, 16–18 m).

*Callechelys kuro* (Kuroda, 1947): McCosker *et al.*, 2011:272.

*Yirrkala kuro* (Kuroda, 1947): Shao *et al.*, 2008:239.

Remarks. A very rare species represented by only a few specimens collected off northeastern and southwestern Taiwan.

***Callechelys marmorata* (Bleeker, 1854)**

雲紋麗鰻

*Dalophis marmorata* Bleeker, 1854:247 (type locality: Sibogha, western Sumatra [Sibolga, Sumatera Utara, Indonesia, eastern Indian Ocean]).

*Callechelys marmorata* (Bleeker, 1854): Shao *et al.*, 2008:238.

Remarks. A widespread species in the Indo-west Pacific Ocean, but only occasionally found in coral reef regions of southern Taiwan.

***Cirrhimuraena chinensis* Kaup, 1856**

中國鬚蛇鰻

*Cirrhimuraena chinensis* Kaup, 1856a:51 (type locality stated as China, Makassar [South Sulawesi, Indonesia] in Kaup, 1856b). Chen & Weng, 1967:59; Chen, 1969:137; Shen 1984a:116; Chen & Yu, 1986:258; Shen *et al.*, 1993:111; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1.

Remarks. An uncommon species collected from southwestern Taiwan and Kinmen Island near China. Although there seems to be more than one species in the western Pacific, our specimens are identical to those collected from Vietnam. This genus requires further taxonomic studies.

**\**Cirricaecula macdowelli* McCosker & Randall, 1993**

麥氏無鰭蛇鰻

*Cirricaecula macdowelli* McCosker & Randall, 1993:190, fig. 1-2 (type locality: Penghu Island, Taiwan strait, 30–50 m). Chen, 2004:30; Ho & Shao, 2011:23, table 1.

Remarks. This species is known only from the holotype collected in shallow water of Penghu Island, southwestern Taiwan.

**\**Echelus polyspondylus* McCosker & Ho, 2015**

多椎鰭蛇鰻

*Echelus polyspondylus* McCosker & Ho, 2015:73, figs. 1–3. (type locality: Dong-gang, southwestern Taiwan).

Remarks. Some specimens of this species were previously misidentified as *Myrophis microchir*, which indicates the similarity between these two groups. This is a deepwater species, occasionally collected by bottom trawl off northeastern and southwestern Taiwan.

***Echelus uropterus* (Temminck & Schlegel, 1846)**

小尾鰭蛇鰻

*Conger uropterus* Temminck & Schlegel, 1846:261 (Type locality: Nagasaki, Japan).

*Echelus uropterus* (Temminck & Schlegel, 1846): Shao *et al.*, 2008:238; Ho *et al.*, 2010:27, table 1.

Remarks. A very common species collected by bottom trawl off northeastern and southwestern Taiwan. The largest specimen exceeds 600 mm TL.

***Lamnostoma mindora* (Jordan & Richardson, 1908)**

龍口舌鰻

*Caecula mindora* Jordan & Richardson, 1908:239, fig. 4 (type locality: Mindoro Island, Philippines).

*Lamnostoma mindora* (Jordan & Richardson, 1908): Hatooka & Yoshino, 1998; Lin, 2007:88.

Remarks. A rare species only found in river mouth areas around Taiwan. Lin (2007) noted that the species can be found in lower reaches of streams in Yilan, Hualien, Taitung and Pingtung.

***Leiuranus semicinctus* (Lay & Bennett, 1839)**

半環平蓋蛇鰻

*Ophisurus semicinctus* Lay & Bennett, 1839:66, pl. 20, fig. 4 (type locality: Oahu? [Hawaiian Islands, central Pacific]).

*Leiuranus semicinctus* (Lay & Bennett, 1839): Chen & Weng, 1967:60; Chen, 1969:137; Shen 1984a:116; Chen & Yu, 1986:258; Shen *et al.*, 1993:111; Shao *et al.*, 2008:238; Ho *et al.*, 2010:table 1.

Remarks. An uncommon species occasionally found in coral reef regions.

***Muraenichthys gymnopterus* (Bleeker, 1853)**

裸鰭蟲鰻

*Muraena gymnopterus* Bleeker, 1853a:52 (type locality: Jakarta, Java, Indonesia).

*Muraenichthys gymnopterus* (Bleeker, 1853a): Chen & Yu, 1986:258.

Remarks. This rare species is only represented by few specimens in the collections of Taiwan.

**+*Muraenichthys hattae* Jordan & Snyder, 1901**

短鰭蟲鰻

*Muraenichthys hattae* Jordan & Snyder, 1901:862, fig. 12 (type locality: rock pool at Wakanoura, Wakayama Prefecture, Japan).

Voucher specimen: NTUM 1075 (Da-xi).

Remarks. A new record, collected in northeastern Taiwan.

**+*Muraenichthys schultzei* Bleeker, 1857**

許氏蟲鰐

*Muraenichthys schultzei* Bleeker, 1857d:366 (type locality: Java, Indonesia, eastern Indian Ocean).

Voucher specimens: USNM 314095, 314099, 314105 (Kenting, S. Taiwan).

Remarks. A new record, collected in southern Taiwan.

**+*Muraenichthys thompsoni* Jordan & Richardson, 1908**

湯氏蟲鰐

*Muraenichthys thompsoni* Jordan & Richardson, 1908:237, fig. 1 (type locality: Manila Bay, the Philippines).

Voucher specimen: NMMB-P 17508 (Dong-gang).

Remarks. A new record, collected in southwestern Taiwan.

**\**Muraenichthys velinasalis* Hibino & Kimura, 2015**

鼻瓣蟲鰐

*Muraenichthys velinasalis* Hibino & Kimura, 2015b:65, figs. 2b, 3f, 4, 5a, b, c (type locality: southwestern Taiwan).

Remarks. A widespread species in the western Pacific Ocean.

***Myrichthys colubrinus* (Boddaert, 1781)**

竹節花蛇鰐

*Muraena colubrina* Boddaert, 1781:56, pl. 2, fig. 3 (type locality: Ambon Island, Molucca Islands, Indonesia).

*Myrichthys colubrinus* (Boddaert, 1781): Chen & Weng, 1967:61; Chen, 1969:137; Shen 1984a:116; Chen & Yu, 1986:258; Shen et al., 1993:112; Shao et al., 2008:238; Ho et al., 2010:table 1.

Remarks. An uncommon species found over sandy bottoms or in sea grass beds near coral reef areas. It is a remarkable mimic of venomous hydrophine sea snakes (McCosker & Rosenblatt, 1993).

***Myrichthys maculosus* (Cuvier, 1816)**

巨斑花蛇鰐

*Muraena maculosa* Cuvier, 1816:232, fig. 29 (type locality: European Seas, probably Indonesia).

*Myrichthys aki* Tanaka, 1917: Chen & Yu, 1986:258; Shen et al., 1993:112.

*Myrichthys maculosus* (Cuvier, 1816): Chen & Yu, 1986:258; Shen et al., 1993:112; Shao et al., 2008:238; Ho et al., 2010:table 1.

Remarks. An uncommon species found in coral reef regions around Taiwan.

**\**Neenchelys cheni* (Chen & Weng, 1967)**

陳氏新鰐

*Myrophis cheni* Chen & Weng, 1967:39 (Type locality: Tungkang [Dong-gang], southwestern Taiwan). Chen, 1969:130; Shen 1984a:117; Chen & Yu, 1986:258; Shen et al., 1993:112; Shao et al., 2008:238.

"*Neenchelys*" *cheni* (Chen & Weng, 1967): Ho et al. 2010:24, table 1; Ho & Shao, 2011:23, table 1.

*Neenchelys cheni* (Chen & Weng, 1967): Hibino et al., 2012:345; Ho et al., 2013:5.

*Neenchelys retropinna* Smith & Böhlke, 1983: McCosker & Chen, 2000:356.

Remarks. An uncommon species collected by midwater trawls off southwestern Taiwan.

**\**Neenchelys diaphora* Ho, McCosker & Smith, 2015**

異新鰻

*Neenchelys diaphora* Ho, McCosker & Smith, 2013:8, fig. 4 (type locality: Dong-gang, southwestern Taiwan). [Name unavailable]

*Neenchelys diaphora* Ho, McCosker & Smith, 2015c:49. (type locality: Dong-gang, southwestern Taiwan).

Remarks. Originally published in Ho *et al.* (2013), but the name is not available because it failed to meet the amended Article 8.5 of the International Code of Zoological Nomenclature (ICZN, 2012). A paper re-naming the species is included in this volume.

**\**Neenchelys gracilis* Ho & Loh, 2015**

*Neenchelys gracilis* Ho & Loh, 2015:54, figs. 1A–B, 2A–B (Type locality: Dong-gang, SW Taiwan).

Remarks. A rare species that is only represented by the holotype.

**\**Neenchelys mccoskeri* Hibino, Ho & Kimura, 2012**

麥氏新鰻

*Muraenichthys macropterus* (not of Bleeker, 1857): Ho *et al.*, 2010:table 1.

*Neenchelys mccoskeri* Hibino, Ho & Kimura, 2012:343 (Type locality: Tungkang, southwestern Taiwan). Ho *et al.*, 2013:9.

Remarks. A deep-sea species commonly collected by bottom trawl off southwestern Taiwan.

***Neenchelys parvipectoralis* Chu, Wu & Jin, 1981**

微鰭新鰻

*Neenchelys parvipectoralis* Chu, Wu & Jin, 1981:24 (type locality: Pingtan Island, Fujian Province, China). Ho *et al.*, 2010:29, table 1; Ho *et al.*, 2013:11.

Remarks. A very common species in the South China Sea, including southwestern Taiwan and Vietnam.

**\**Neenchelys pelagica* Ho, McCosker & Smith, 2015**

大洋新鰻

*Neenchelys pelagica* Ho, McCosker & Smith, 2013: 14, fig. 8 (type locality: Tungkang, southwestern Taiwan). [Name unavailable]

*Neenchelys pelagica* Ho, McCosker & Smith 2015c:50. (type locality: Tungkang, southwestern Taiwan).

Remarks. Originally published in Ho *et al.* (2013), but the name is not available because it failed to meet the amended Article 8.5 of the International Code of Zoological Nomenclature (ICZN, 2012). A paper re-naming the species is included in this volume.

**\**Neenchelys similis* Ho, McCosker & Smith, 2015**

長身新鰻

*Neenchelys deadalus* (non McCosker, 1982). McCosker, 1982:65 (in part).

*Neenchelys similis* Ho, McCosker & Smith, 2013: 15, fig. 9 (Type locality: Tashi, northeastern Taiwan). [Name unavailable]

*Neenchelys similis* Ho, McCosker & Smith, 2015c:50. (Type locality: Tashi, northeastern Taiwan).

Remarks. Originally published in Ho *et al.* (2013), but the name was not available because it failed to meet the

amended Article 8.5 of the International Code of Zoological Nomenclature (ICZN, 2012). A paper re-naming the species is included in this volume. A rare species only represented by one specimen collected from northeastern Taiwan.

***Ophichthus altipennis* (Kaup, 1856)**

帆鰭蛇鰻

*Microdonophis altipennis* Kaup, 1856a:43 (type locality: Makassar, Sulawesi, Indonesia).

*Pisoodonophis zophistius* Jordan & Snyder, 1901: Chen et al., 2010:28.

*Ophichthus altipennis* (Kaup, 1856). McCosker & Ho, 2015:73.

Remarks. Chen et al. (2010) firstly provided underwater photographs identified as *Pisoodonophis zophistius*, which is a junior synonym of this species. McCosker & Randall (2002) discussed the nomenclatural confusion about this species and established the validity of *Ophichthus altipennis*. Two voucher specimens were found (NTUM 1793 and NTUM 50567) in the collection by YH.

**\**Ophichthus aphotistos* McCosker & Chen, 2000**

深海黑體蛇鰻

*Ophichthus aphotistos* McCosker & Chen, 2000:354, fig. 1 (type locality: Tung-Kong [Tongkang] Channel, southwestern Taiwan, 22°22'N, 120°19'E, 700–800 m). Shao et al., 2008:238; Ho & Shao, 2011:23, table 1; McCosker & Ho, 2015:72.

Remarks. A rare species represented only by the type series and one additional specimen (ASIZP 64278), collected in northeastern Taiwan.

***Ophichthus apicalis* (Bennett, 1830)**

頂蛇鰻

*Ophisurus apicalis* Bennett, 1830:692 (type locality: Sumatra, Indonesia).

*Ophichthus apicalis* (Bennett, 1830): ?Shen et al., 1993:113; ?Shao et al., 2008:238; ?Ho et al., 2010:table 1; ?Chiu et al. 2013:206; McCosker & Ho, 2015:73.

*Ophichthys apicalis* (Bennett, 1830): ?Chen & Weng, 1967:72; ?Chen, 1969:137; ?Shen 1984a:118; ?Chen & Yu, 1986:259.

Remarks. This name has been applied to more than one species. Several recently described species, including *O. machidai*, *O. obtusus* and *O. shaoi* from Taiwan have all previously been misidentified as this species. As the type was lost, no details are available for this species. Chiu et al. (2013) recognized this species but without providing any voucher for it. McCosker & Ho (2015) recognized three specimens as the present species.

***Ophichthus asakusae* Jordan & Snyder, 1901**

淺草氏蛇鰻

*Ophichthus asakusae* Jordan & Snyder, 1901:872, fig. 18 (Type locality Bay of Tokyo, near Misaki, Japan). Chiu et al., 2013:206; McCosker & Ho, 2015:73.

Remarks. Chiu et al. (2013) included this species in their key but did not provide a voucher. It has only been collected from a few localities in Taiwan; most specimens collected from shallow water off Ke-tzu-liao, southwestern Taiwan.

**\**Ophichthus bicolor* McCosker & Ho, 2015**

雙色蛇鰻

*Ophichthus bicolor* McCosker & Ho, 2015:76, figs. 4–6 (type locality: Dong-gang, southwestern Taiwan).

Remarks. A deepwater species occasionally collected from southeastern Taiwan by hook and line and southwestern Taiwan by bottom trawl.

***Ophichthus bonaparti* (Kaup, 1856)**

鮑氏蛇鰻

*Poecilocephalus bonaparti* Kaup, 1856a:43 (type locality: Ambon Island, Molucca Islands, Indonesia).  
*Ophichthus bonaparti* (Kaup, 1856): Chen, 1993:20; Chen, 2004:30; McCosker & Ho, 2015:72.

Remarks. Occasionally photographed by SCUBA divers in southern Taiwan. Chen (1993, 2004) recorded this species from the Penghu Islands.

***Ophichthus cephalozona* (Bleeker, 1864)**

頸斑蛇鰻

*Ophichthus cephalozona* Bleeker, 1864c:49, pl. 156, fig. 2 (type locality: Singapore; Ambon Island, Molucca Islands, Indonesia). ?Shen, 1988:15; ?Shen et al., 1993:113; Chen, 2004:30; McCosker & Ho, 2015:72.  
*Ophichthys cephalozona* (Bleeker, 1864c): ?Jordan & Evermann, 1902:326; ?Matsubara, 1955:349; ?Chen, 1956:76; ?Chen & Weng, 1967:69; ?Chen, 1969:137; ?Shen 1984a:117; ?Chen & Yu, 1986:259.

Remarks. Although most records have been determined to be misidentifications of *O. lithinus*, an underwater photograph examined by one of us (JMC) confirms the presence of this species in Taiwan.

***Ophichthus erabo* (Jordan & Snyder, 1901)**

斑紋蛇鰻

*Microdonophis erabo* Jordan & Snyder, 1901:870, fig. 17 (type locality: Misaki, Japan). Chen & Weng, 1967:67; Chen, 1969:137; Chen & Yu, 1986:259; Shen et al., 1993:111.  
*Ophichthus erabo* (Jordan & Snyder, 1901): Shen, 1988:15; ?Chen, 2004:30; Shao et al., 2008:238; Chiu et al., 2013:206; McCosker & Ho, 2015:72.  
*Ophichthys erabo* (Jordan & Snyder, 1901): Shen 1984a:117.

Remarks. A common species collected by bottom trawl off northeastern and southwestern Taiwan; more abundant in southwestern Taiwan.

***Ophichthus fasciatus* (Chu, Wu & Jin, 1981)**

條紋蛇鰻

*Microdonophis fasciatus* Chu, Wu & Jin, 1981:21 (type locality: Pingtan Island, Fujian Province, China). Chen, 1993:19; Chen, 2004:30.  
*Ophichthus fasciatus* (Chu, et al., 1981): McCosker & Ho, 2015:72.

Remarks. First recorded by Chen (1993) from Penghu. The voucher has not been examined and might have been lost.

***Ophichthus lithinus* (Jordan & Richardson, 1908)**

艾氏蛇鰻

*Leiuranus lithinus* Jordan & Richardson, 1908:238, fig. 3 (type locality: Cuyo, Philippines)  
*Ophichthus lithinus* (Jordan & Richardson, 1908): Ho et al., 2010:table 1; Ho & Shao, 2011:23, table 1; McCosker & Ho, 2015:72.  
*Ophichthus evermanni* Jordan & Richardson, 1909:172 (Type locality: Takao [Kaohsiung], Taiwan). Jordan et al., 1913:87; Chen, 1956:76; Chen & Weng, 1967:70; Chen, 1969:137; Shen 1984a:117; Chen & Yu, 1986:259; Shen et al., 1993:113; Shao et al., 1994a:276; Chen, 2004:30.  
*Ophichthys cephalozona* (not of Bleeker, 1864c): Jordan & Evermann, 1902:326.

Remarks. A common species collected from shallow waters around Taiwan, especially in the southwest.

***Ophichthus machidai* McCosker, Ide & Endo 2012**

町田氏蛇鰐

*Ophichthus machidai* McCosker, Ide & Endo 2012:8, figs. 4–6 (type locality: Tosa Bay, Saga fish market (33°04.48'N, 133°06.80'E), Kuroshio-chu, Kochi Prefecture, Japan). Chiu *et al.*, 2013:204; McCosker & Ho, 2015:73.

Remarks. Previous records of *O. apicalis* may include misidentification of this species. It is common in bycatch of bottom trawls.

***Ophichthus macrochir* (Bleeker, 1853)**

大鰭蛇鰐

*Ophisurus macrochir* Bleeker, 1853a:26 (type locality: Jakarta, Java, Indonesia).

*Ophichthus macrochir* (Bleeker, 1853a): Shen *et al.*, 1993:113; Shao *et al.*, 2008:238; Chen, 2004:30; Ho *et al.*, 2010:19, table 1; McCosker & Ho, 2015:72.

*Ophichthys macrochir* (Bleeker, 1853a): Chen & Weng, 1967:73; Chen, 1969:137; Shen 1984a:118; Chen & Yu, 1986:259.

Remarks. Most specimens examined have 207–218 total vertebrae. However, one specimen with 186 total vertebrae is possibly a different species or has a damaged tail.

**+*Ophichthus megalops* Asano, 1987**

大眼蛇鰐

*Ophichthus megalops* Asano, 1987:135, figs. 1–2 (type locality: Owase, Mie Prefecture, Japan, ca. 360 m). ?Ho *et al.*, 2010:table 1; McCosker & Ho, 2015:72.

Voucher specimens: NMMB-P12006, 12193 (Nan-fang-ao); NMMB-P17751 (Daxi).

Remarks. New record for Taiwan. Although Ho *et al.* (2010) identified a specimen as this species, that specimen is now recognized as an unidentified species. The specimens on which this record is based were collected in northeastern Taiwan.

***Ophichthus obtusus* McCosker, Ide & Endo, 2012**

壯體蛇鰐

*Ophichthus obtusus* McCosker, Ide & Endo 2012:12, figs. 7–9 (type locality: Tosa Bay, Irino fish market, Kuroshio-cho, Kochi Prefecture, Japan). Chiu *et al.*, 2013:204; McCosker & Ho, 2015:73.

Remarks. A common species collected by bottom trawl around Taiwan.

***Ophichthus polyophthalmus* (Bleeker, 1864)**

眼斑蛇鰐

*Ophichthys polyophthalmus* Bleeker, 1864b:43 (type locality: Amboin Island, Molucca Islands, Indonesia). Shen 1984a:117.

*Ophichthus polyophthalmus* (Bleeker, 1864b): Chen, 2004:31; McCosker & Ho, 2015:72.

*Microdonophis polyophthalmus* (Bleeker, 1864b): Okada, 1938:152; Chen, 1956:76; Chen & Weng, 1967:67; Chen, 1969:137; Chen & Yu, 1986:259; Shen *et al.*, 1993:112.

Remarks. Although recorded by many authors, there is only one specimen present in FRIP with confirmed identification.

**\**Ophichthus shaoi* McCosker & Ho, 2015**

邵氏蛇鰐

*Ophichthus shaoi* McCosker & Ho, 2015:80. figs. 7–9 (type locality: Da-xi, Yilan, northeastern Taiwan).

Remarks. A species occasionally collected by bottom trawl in northeastern and southwestern Taiwan.

***Ophichthus urolophus* (Temminck & Schlegel, 1846)**

裙鰭蛇鰐

*Conger urolophus* Temminck & Schlegel, 1846:260, pl. 114 (type locality: Nagasaki, Japan).

*Ophichthus tsuchidae* (Jordan & Snyder, 1901): Shen et al., 1993:114.

*Ophichthus urolophus* (Temminck & Schlegel, 1846): Shen et al., 1993:114; Shao et al., 2008:238; Ho et al., 2010:table 1; Chiu et al., 2013:206; McCosker & Ho, 2015:73.

*Ophichthys urolophus* (Temminck & Schlegel, 1846): Chen & Weng, 1967:71; Chen, 1969:137; Shen 1984a:117; Chen & Yu, 1986:259.

*Ophichthys tsuchidai* (Jordan & Snyder, 1901): Chen & Yu, 1986:259; Shao et al., 2008:238; Chiu et al., 2013:206.

*Microdonophis intermedius* (Regan, 1905): Liang, 1951:11; Chen, 1956:76.

Remarks. This is a very common species collected by bottom trawl from shallow to deep waters around Taiwan. In our examination of numerous specimens, we noted that the number of tooth rows changes with growth. There is usually a single row of teeth on both maxilla and dentary in specimens less than 300 mm TL, becoming biserial in specimens larger than 300 mm TL and triserial in specimens larger than 600 mm TL. Several authors have recognized *O. tsuchidae* from Taiwan, which is a junior synonym of *O. urolophus*.

**+*Ophisurus macrorhynchos* Bleeker, 1853**

巨吻沙蛇鰐

*Ophisurus macrorhynchos* Bleeker, 1853a:28 (type locality: near Kaminoseki Island, Yamaguchi Prefecture, Japan).

Voucher specimens: ASIZP 62582, 64629 (Da-xi)

Remarks. A new record for Taiwan; a rare species occasionally collected in deep water by bottom trawl off northeastern Taiwan. The largest individual is more than 2 meters long (Ho, pers. obser.).

***Phyllophichthus xenodontus* Gosline, 1951**

葉鼻鰐

*Phyllophichthus xenodontus* Gosline, 1951:316 (type locality: Hauula Park, Oahu Island, Hawaiian Islands): Ho et al., 2010:30, table 1.

Remarks. The species was recorded by Ho et al. (2010) from two specimens collected from southwestern Taiwan. There is also a specimen (USNM 314694) collected in 1968. It is a rare species that is not commonly seen in Taiwan.

***Pisodonophis boro* (Hamilton, 1822)**

波路荳齒蛇鰐

*Ophisurus boro* Hamilton, 1822:20 (type locality: Ganges estuary near Calcutta, India).

*Pisodonophis boro* (Hamilton, 1822): Chen, 1956:76; Chen & Weng, 1967:63; Chen, 1969:137; Shen 1984a:117; Chen & Yu, 1986:259; Shen et al., 1993:114; Shao et al., 2008:238; Ho et al., 2010:table 1.

Remarks. A common species found in the bycatch of bottom trawls off southwestern Taiwan, but it is less abundant than *P. cancrivorus*.

***Pisodonophis cancrivorus* (Richardson, 1848)**

食蟹壹齒蛇鰻

*Ophisurus cancrivorus* Richardson, 1848b:97 (type locality: Port Essington, Northern Territory, Australia).

*Pisodonophis cancrivorus* (Richardson, 1848): Jordan & Evermann, 1902:326; Jordan & Richardson, 1909:172; Chen, 1956:76; Chen & Weng, 1967:65; Lee, 1967:44; Chen, 1969:137; Shen 1984a:116; Chen & Yu, 1986:258; Shen, 1988:15; Shen et al., 1993:114; Chen & Fang, 1999:54; Shao et al., 2008:238; Chen, 2004:31; Ho et al., 2010:table 1.

Remarks. A very common and abundant species found in shallow waters around Taiwan, especially over mud bottoms.

**\**Pylorobranchus* McCosker & Chen, 2012**

守鰓鰻屬

*Pylorobranchus* McCosker & Chen in McCosker et al., 2012:1189 (Type species: *Pylorobranchus hoi*, by monotypy). McCosker, 2014:334

**\**Pylorobranchus hoi* McCosker, Loh & Lin, 2012**

何氏守鰓鰻

*Pylorobranchus hoi* McCosker, Loh & Lin in McCosker et al., 2012:1192, figs. 1–4 (type locality: Changbin, 23°18'N, 121°29'E, Taitung County, Taiwan, 300 m). McCosker, 2014:337.

Remarks. A deepwater species known only from the type series collected by hook and line off southeastern Taiwan.

**+*Schultzidium johnstonensis* (Schultz & Woods, 1949)**

舒氏鰻

*Muraenichthys johnstonensis* Schultz & Woods, 1949:172, fig. 1 (type locality: Johnston Island [Central Pacific]).

Voucher specimen: NTUM 10694 (Kin-men, Taiwan Strait).

Remarks. This is a new record for Taiwan, and is represented by a single specimen collected at Kin-men, western Taiwan Strait.

**+*Scolecenchelys fuscovenensis* McCosker, Ide & Endo, 2012**

黑尾蟲鰻

*Scolecenchelys fuscovenensis* McCosker, Ide & Endo, 2012:2, figs. 1–3 (type locality: Tosa Bay, 33°14.70'N, 133°38.42'E - 33°13.01'N, 133°36.55'E, Japan, 223–269 m).

Voucher specimens: ASIZP 53979, NMMB-P 12004 (Dong-gang).

Remarks. This is a new record for Taiwan. Both vouchers were collected by bottom trawl off southwestern Taiwan.

**+*Scolecenchelys iredalei* (Whitley, 1927)**

愛氏蟲鰻

*Muraenichthys iredalei* Whitley, 1927:5, fig. 1 (type locality: Michaelmas Cay, Great Barrier Reef, off Cairns, Queensland, Australia). Hibino and Kimura, 2015a:unnumbered.

Voucher specimen: USNM 427603 (Pingtung).

Remarks. This is a new record for Taiwan. Although McCosker et al. (2006) treated this as a junior synonym of *S. australis*, Hibino & Kimura (2015a) recognized it as a distinct species. The voucher was collected from a shallow coral reef of southern Taiwan.

**+*Scolecenchelys laticaudata* (Ogilby, 1897)**

側尾蟲鰻

*Myropterura laticaudata* Ogilby, 1897:247 (type locality: Fiji Islands [South Pacific]).

*Scolecenchelys gymnota* (non Bleeker, 1854): Shao *et al.*, 2008:239.

*Scolecenchelys laticaudata* (Ogilby, 1897): Hibino & Kimura, 2015a:unnumbered.

Voucher specimens: ASIZP 55133 (Hengchun, Pingtung); USNM 313981 (Kenting, Pingtung).

Remarks. This is a new record for Taiwan. Specimens previously referred to *Scolecenchelys gymnota* (ASZIP 55133) are identified as this species (Hibino & Kimura, 2015a). The USNM specimen was collected from a shallow coral reef in southern Taiwan.

**+*Scolecenchelys macroptera* (Bleeker, 1857)**

大鰭蟲鰻

*Muraenichthys macropterus* Bleeker, 1857a:91 (type locality: Ambon Island, Molucca Islands, Indonesia).

Voucher specimen: USNM 314037 (Hengchun, Pingtung).

Remarks. This is a new record for Taiwan. The voucher specimen was collected from a tide pool in southern Taiwan.

***Skythrenchelys zebra* Castle & McCosker, 1999**

龜蛇鰻

*Skythrenchelys zebra* Castle & McCosker, 1999:116, figs. 1A–C, 2A (type locality: Thevara, Ernakulam, southern India, 10°00'N, 76°16'E, shallow turbid water). Chiu *et al.*, 2013:202.

Remarks. A rare species represented by only a few specimens collected from southwestern Taiwan by bottom trawl. Other specimens collected from China and Vietnam were examined by HH.

**\**Sympenchelys* Hibino, Ho & Kimura, 2015**

扁鰻屬

*Sympenchelys* Hibino, Ho & Kimura, 2015:42. (type species: *Sympenchelys taiwanensis* Hibino, Ho & Kimura, by monotypy).

**\**Sympenchelys taiwanensis* Hibino, Ho & Kimura, 2015**

台灣扁鰻

*Sympenchelys taiwanensis* Hibino, Ho & Kimura: 2015:44. figs. 1–6 (type locality: Dong-gang, southwestern Taiwan, South China Sea, ca. 200 m).

Remarks. A rare and small species occasionally collected by bottom trawl off southwestern Taiwan.

**\**Xyrias chioui* McCosker, Chen & Chen, 2009**

邱氏列齒鰻

*Xyrias chioui* McCosker, Chen & Chen, 2009:63 (type locality: Changbin, 23°17'N, 121°27'E, Taitung County, Taiwan, 60–70 m). Ho & Shao, 2011:23, table 1.

Remarks. A rare species known only from the holotype collected by hook and line off southeastern Taiwan.

**+*Xyrias revulsus* Jordan & Snyder, 1901**

列齒鰻

*Xyrias revulsus* Jordan & Snyder, 1901:869, fig. 16 (Type locality: Misaki, Japan).

Voucher specimens: ASIZP 70172 (Da-xi); NMMB-P16531, 16532 (Dong-gang).

Remarks. Although Tan (1983) recorded this species from Diao-yu Island, specimens of this species were recently collected and represent the first confirmed record from Taiwan. It is a relatively deepwater species collected by bottom trawl off northeastern and southwestern Taiwan.

**+*Yirrkala misolensis* (Günther, 1872)**

米蘇爾細鋤蛇鰻

*Ophichthys misolensis* Günther, 1872b:426 (type locality: Misol Island, Irian Jaya, Indonesia).

Voucher specimens: NMMB-P 12003 (Dong-gang); NMMB-P 17506 (Dong-gang).

Remarks. This is a new record for Taiwan; represented by only a few specimens collected by bottom trawl in southwestern Taiwan. It is very common in Vietnam, e.g. Nha Trang and Mui Ne, collected from depths less than 100 meters, based on HH's observation.

**Family Serrivomeridae**

鋸犁鰻科

***Serrivomer sector* Garman, 1899**

長齒鋸犁鰻

*Serrivomer sector* Garman, 1899:320, pl. 63 (type locality: Eastern Pacific, 3°–7°N, 79°–86°W, 0–1772 fathoms [0–3240 m]).  
Shao et al., 2008:239.

Remarks. This rare species is represented by only one specimen collected from Taiwan.

**Family Synaphobranchidae**

合鰓鰻科

***Dysomma anguillare* Barnard, 1923**

前肛鰻

*Dysomma anguillaris* Barnard, 1923:443 (type locality: KwaZulu-Natal, South Africa, southwestern Indian Ocean, 63 fathoms [115 m]). Chen & Weng, 1967:83; Chen, 1969:138; Shen 1984a:110; Chen & Yu, 1986:260; Shen et al., 1993:109.  
*Dysomma anguillare* Barnard, 1923: Chen & Mok, 2001:79; Shao et al., 2008:238; Ho et al., 2010:table 1; Ho et al., 2015d:91.  
?*Dysomma angustum*: Chen, 1956:73.

Remarks. A very common species collected by bottom trawl and midwater trawl around Taiwan. It is far more abundant than all other congeners in Taiwan.

***Dysomma dolichosomatum* Karrer, 1983**

長身前肛鰻

*Dysomma dolichosomatum* Karrer, 1983:93, figs. 28 A–B (type locality: Madagascar, 22°25'S, 43°04.5'E, 550–555 m). Shen et al., 1993:109; Chen & Mok, 2001:79; Shao et al., 2008:238; Ho et al., 2015d:93.

Remarks. A common species collected by midwater trawl off northeastern and southwestern Taiwan. It usually feeds on midwater shrimps.

***Dysomma goslinei* Robins & Robins, 1976**

高氏前肛鰻

*Dysomma goslinei* Robins & Robins, 1976:261, figs. 3, 6, 7c, 8 (type locality: Indian Ocean, 14°52'N, 96°39'E). Shen et al., 1993:109; Chen & Mok, 2001:79; Ho et al., 2015d:94.

Remarks. A rare species represented by several specimens in Taiwan.

**\**Dysomma longirostrum* Chen & Mok, 2001**

長吻前肛鰻

*Dysomma longirostrum* Chen & Mok, 2001:79, figs. 1A, 2 (type locality: Nanfangao fish market, northeastern Taiwan, 100–150 m). Shao *et al.*, 2008:238; Ho & Shao, 2011:22, table 1; Ho *et al.*, 2015d:95.

Remarks. A rare species represented by several specimens in collections worldwide. In Taiwan it is represented only by the holotype.

**\**Dysomma melanurum* Chen & Weng, 1967**

黑尾前肛鰻

*Dysomma melanurum* Chen & Weng, 1967:84, fig. 63 (type locality: Tungkong [Dong-gang], southwestern Taiwan). Chen, 1969:138; Shen, 1984a:110; Robins & Robins in Böhlke, 1989:249; Shen *et al.*, 1993:109; Smith, 1999a:1661; Chen & Mok, 2001:79; Shao *et al.*, 2008:238; Ho *et al.*, 2010:26, table 1; Ho & Shao, 2011:22, table 1; Ho *et al.*, 2015d:96.

Remarks. This species is occasionally collected by midwater trawl from southwestern Taiwan; feeds on midwater shrimps.

**\**Dysomma opisthoproctus* Chen & Mok, 1995**

後臀前肛鰻

*Dysomma opisthoproctus* Chen & Mok, 1995:927, fig. 1 (type locality: off the coast of Nan-Fang-Ao, northeastern Taiwan, 200 m). Smith, 1999a:1661; Chen & Mok, 2001:79; Shao *et al.*, 2008:238; Ho & Shao, 2011:22, table 1; Ho *et al.*, 2015d:98.

Remarks. A rare species, to date known only from the holotype.

***Dysomma polycatodon* Karrer, 1983**

多齒前肛鰻

*Dysomma polycatodon* Karrer, 1983:89, figs. 27 A–B (type locality: Madagascar, 15°21'S, 46°11'E, 170–175 m). Chen & Mok, 2001:79; Ho *et al.*, 2015d:99.

Remarks. A common species collected by bottom trawl from northeastern and southwestern Taiwan, but not as abundant as *D. anguillare*.

**\**Dysomma taiwanensis* Ho, Smith & Tighe, 2015**

台灣前肛鰻

*Dysomma taiwanensis* Ho, Smith & Tighe, 2015d:87, figs. 1A–B, 2A–B (type locality: Daxi, Yilan, northeastern Taiwan).

Remarks. A rare deepwater species represented only by the type series collected by bottom trawl off southeastern Taiwan.

***Dysommina rugosa* Ginsburg, 1951**

多皺短身前肛鰻

*Dysommina rugosa* Ginsburg, 1951:450, fig. 6 (type locality: off Cumberland Island, Georgia, U.S.A., western North Atlantic, 30°53'00"N, 79°42'30"W, 273 fathoms [499 m]). Chen & Mok, 2001:79; Shao *et al.*, 2008:238; Ho *et al.*, 2015d:100.

Remarks. A rare species occasionally collected by bottom trawl from northeastern and southwestern Taiwan.

**+*Histiobranchus bathybius* (Günther, 1877)**

深海旗鰓鰻

*Synaphobranchus bathybius* Günther, 1877:445 (type locality: south of Tokyo, Japan and South Atlantic between Cape of Good Hope and Kerguelen Island).

Voucher specimen: ASIZP 66208 (Suao, Yilan, northeastern Tawian).

Remarks. This is a new record for Taiwan, and it is only represented by a single specimen collected from northeastern Taiwan at 2204–2334 m.

***Ilyophis brunneus* Gilbert, 1891**

褐泥鰻

*Ilyophis brunneus* Gilbert, 1891:352 (type locality: San Cristóbal Island [Chatham Island], Galápagos Islands, 0°36'30"S, 89°19'00"W, 634 fathoms [1159 m]). Chen & Mok, 2001:79; Shao *et al.*, 2008:238.

Remarks. A rare species represented only by a few specimens collected by bottom trawl off northeastern Tawian.

***Meadia abyssalis* (Kamohara, 1938)**

箭齒前肛鰻

*Dysomma abyssale* Kamohara, 1938:12, fig. 3 (type locality: Sagami Sea, Japan).

*Meadia abyssalis* (Kamohara, 1938): Chen & Mok, 2001:79; Shao *et al.*, 2008:238.

Remarks. A deepwater species commonly collected by hook and line from eastern Taiwan and occasionally collected by bottom trawl off northeastern and southwestern Taiwan.

**\**Meadia roseni* Mok, Lee & Chen, 1991**

羅氏箭齒前肛鰻

*Meadia roseni* Mok, Lee & Chan, 1991:39, figs. 1–3 (type locality: 22°21'05"N, 120°12'46"E. [off southwestern Taiwan], 1020 m). Shen *et al.*, 1993:109; Smith, 1999a:1661; Chen & Mok, 2001:79; Shao *et al.*, 2008:238; Ho & Shao, 2011:22, table 1.

Remarks. A species commonly collected from deep waters by bottom trawl off southwestern Taiwan.

***Simenchelys parasitica* Gill, 1879**

寄生鰻

*Simenchelys parasiticus* Gill in Goode & Bean, 1879:27 (type locality: Banquereau, Canada). Shen *et al.*, 1993:110; Chen & Mok, 2001:79.

Remarks. A rare species only represented by several specimens collected from deep waters by bottom trawl off northeastern Taiwan.

***Synaphobranchus affinis* Günther, 1877**

合鰓鰻

*Synaphobranchus affinis* Günther, 1877:445 (type locality: Japan). Shen *et al.*, 1993:110; Chen & Mok, 2001:79; Shao *et al.*, 2008:238.

Remarks. There are many specimens deposited in the ASIZP collection. However, there are taxonomic problems remaining, and this group needs further study.

***Synaphobranchus brevidorsalis* Günther, 1887**

短背鰭合鰓鰻

*Synaphobranchus brevidorsalis* Günther, 1887:255 (type locality: North of New Guinea and Japan). Shao *et al.*, 2008:238.

Remarks. There are several specimens of this species in the ASIZP collection, most of them collected from the South China Sea off Taiwan.

***Synaphobranchus kaupii* Johnson, 1862**

高氏合鰓鰻

*Synaphobranchus kaupii* Johnson, 1862:169 (type locality: off Madeira, eastern Atlantic). Chen & Mok, 2001:79; Shao *et al.*, 2008:238.

Remarks. Although there are many specimens found in the collection, taxonomic problems remain, and this group needs further study.

**Order Saccopharyngiformes**

寬咽鰻目

**+Family Eurypharyngidae**

寬咽鰻科

**+*Eurypharynx pelecanoides* Vaillant, 1882**

寬咽鰻

*Eurypharynx pelecanoides* Vaillant, 1882:1226 (type locality: off Morocco, 2300 m).

Voucher specimens: ASIZP 60920, 62122, 62597 (Daxi).

Remarks. This is a new record in Taiwan, also a rare species occasionally collected by the bottom trawl in Taiwan.

**Acknowledgements**

Many individuals are acknowledged for their assistance, including P.-L. Lin (ASIZP), C.-G. Zhang (ASIZB), M. Hong, J. Fong (CAS), J. Maclaine, O. Crimmen (BMNH), J. Pogonoski (CSIRO), S. Kimura (FRLM), Z. Gabsi, C. Ferrara, R. Causse, P. Pruvost (MNHN), G. Shinohara, M. Nakae (NSMT), W.-J. Chen (NTUM), S. Raredon, J. Williams, J. Clayton, D. Pitassy, S. Smith, K. Murphy (USNM). This project is supported by the National Museum of Marine Biology & Aquarium.

**References**

- Abbott, C.C. (1860) Description of new species of apodal fishes in the museum of the Academy of Natural Sciences of Philadelphia. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 12, 475–479.
- Abe, T., Miki, M. & Asai, M. (1977) Description of a new garden eel from Japan. *Uo (Japanese Society of Ichthyologists)*, 28, 1–8.
- Ahl, J.N. (1789) *Specimen ichthyologicum de Muraena et Ophichtho, quod venia exp. fac. Med. ups. Praeside Carol. Pet. Thunberg, etc. Upsala*, 14 pp.
- Alcock, A.W. (1889) Natural history notes from H. M. Indian marine survey steamer 'Investigator,' Commander Alfred Carpenter, R. N., D. S. O., commanding.--No. 13. On the bathybiial fishes of the Bay of Bengal and neighbouring waters,

- obtained during the seasons 1885-1889. *Annals and Magazine of Natural History (Ser. 6)*, 4, 450-461.
- Asano, H. (1958) Studies on the conger eels of Japan. I. Description of two new subspecies referable to the genus *Allociongaster*. *Dobutsugaku Zasshi [Zoological Magazine Tokyo]*, 67 (7), 191-196. [In Japanese, English summary.]
- Asano, H. (1962) Studies on the congrid eels of Japan. *Bulletin of the Misaki Marine Biological Institute Kyoto University*, 1, 1-143.
- Asano, H. (1987) A new ophichthid eel, *Ophichthus megalops*, from the Kumano-nada, Japan. *Japanese Journal of Ichthyology*, 34 (2), 135-137.
- Barnard, K.H. (1923) Diagnoses of new species of marine fishes from South African waters. *Annals of the South African Museum*, 13 (pt. 8, no.14), 439-445.
- Beebe, W. & Tee Van, J. (1933) Nomenclatural notes on the shore fishes of Bermuda. *Zoologica (N. Y.)*, 13 (7), 133-158.
- Bennett, E.T. (1831) Observations on a collection of fishes from the Mauritius, presented by Mr. Telfair, with characters of new genera and species. *Proceedings of the Committee of Science and Correspondence of the Zoological Society of London 1830-31 (pt 1)*, 126-128.
- Bennett, E.T. (1830) Class Pisces. In: *Memoir of the life and public services of Sir Thomas Stamford Raffles, F.R.S. &c., Particularly in the government of Java, 1811-1816, and of Bencoolen and its dependencies, 1817-1824*; with details of the commerce and resources of the Eastern archipelago, and selections from his correspondence, by his Widow [Lady Stamford Raffles]. John Murray, London, pp. 686-694.
- Bennett, E.T. (1832) Observations on a collection of fishes from the Mauritius, presented by Mr. Telfair, with characters of new genera and species. *Proceedings of the Committee of Science and Correspondence of the Zoological Society of London, 1830-31 (pt. 1)*, 165-169.
- Bennett, E.T. (1833) Characters of new species from the Mauritius. *Proceedings of the Zoological Society of London*, 1833 (pt 1), 1-32.
- Bleeker, P. (1852) Bijdrage tot de kennis der ichthyologische fauna van de Molucksche Eilanden. Visschen van Amboina en Ceram. *Natuurkundig Tijdschrift voor Nederlandsch Indië*, 3, 229-309.
- Bleeker, P. (1853a) Bijdrage tot de kennis der Muraeoïden en Symbranchoïden van den Indischen Archipel. *Verhandelingen van het Bataviaasch Genootschap van Kunsten en Wetenschappen*, 25 (art. 5), 1-62.
- Bleeker, P. (1853b) Derde bijdrage tot de kennis der ichthyologische fauna van Celebes. *Natuurkundig Tijdschrift voor Nederlandsch Indië*, 3, 739-782.
- Bleeker, P. (1853c) Diagnostische beschrijvingen van nieuwe of weinig bekende vischsoorten van Sumatra. Tiental V-X. *Natuurkundig Tijdschrift voor Nederlandsch Indië*, 4, 243-302.
- Bleeker, P. (1854) Derde bijdrage tot de kennis der ichthyologische fauna van Ceram. *Natuurkundig Tijdschrift voor Nederlandsch Indië*, 5, 233-248.
- Bleeker, P. (1857a) Achtste bijdrage tot de kennis der vischfauna van Amboina. *Acta Societatis Regiae Scientiarum Indo-Néerlandicae*, 2 (7), 1-102.
- Bleeker, P. (1857b) Tweede bijdrage tot de kennis der ichthyologische fauna van Boero. *Natuurkundig Tijdschrift voor Nederlandsch Indië*, 13, 55-82.
- Bleeker, P. (1857c) *Vierde bijdrage tot de kennis der ichthyologische fauna van Japan*. 3 (art. 10), 1-46. [plates. 1-4]
- Bleeker, P. (1857d) Descriptiones specierum piscium javanensium novarum vel minus cognitarum diagnosticae. *Natuurkundig Tijdschrift voor Nederlandsch Indië*, 13, 323-368.
- Bleeker, P. (1858a) Vierde bijdrage tot de kennis der ichthyologische fauna van Timor. Visschen van Atapoepoe. *Natuurkundig Tijdschrift voor Nederlandsch Indië*, 17, 129-140.
- Bleeker, P. (1858b) Vijfde bijdrage tot de kennis der ichthyologische fauna van Japan. *Acta Societatis Regiae Scientiarum Indo-Néerlandicae*, 5, 1-12.
- Bleeker, P. (1859) Over eenige vischsoorten van de Zuidkust-wateren van Java. *Natuurkundig Tijdschrift voor Nederlandsch Indie*, 19, 329-352.
- Bleeker, P. (1863) Notice sur deux espèces inédites du genre *Aphthalmichthys* Kp. *Nederlandsch Tijdschrift voor de Dierkunde*, 1, 163-166.
- Bleeker, P. (1864a) Notice sur quelques poissons de la Baie de Manille. *Nederlandsch Tijdschrift voor de Dierkunde*, 2, 30-32.
- Bleeker, P. (1864b) Poissons inédits indo-archipelagiques de l'ordre des Murènes. *Nederlandsch Tijdschrift voor de Dierkunde*, 2, 38-54.
- Bleeker, P. (1864c) *Atlas ichthyologique des Indes Orientales Néerlandaises, publié sous les auspices du Gouvernement colonial néerlandais*. Tome IV. Murènes, Synbranches, Leptocéphales, 4, 1-150.
- Bleeker, P. (1856) Verslag omtrent eenige vischsoorten gevangen aan de Zuidkust van Malang in Oost-Java. *Natuurkundig Tijdschrift voor Nederlandsch Indië*, 11, 81-92.
- Bleeker, P. (1879) Énumération des espèces de poissons actuellement connues du Japon et description de trois espèces inédites. *Verhandelingen der Koninklijke Akademie van Wetenschappen, Afdeeling Natuurkunde (Amsterdam)*, 18, 1-33. [plates. 1-3]
- Bliss, R. (1883) Descriptions of new species of Mauritian fishes. *Transactions de la Société Royale des Arts et de Sciences de Maurice (N. S.)*, 13, 45-63.
- Bloch, M.E. & Schneider, J.G. (1801) *M. E. Blochii, Systema Ichthyologiae Iconibus CX Illustratum*. Berolini. Ix + 584 pp. [110 plates]

- Boddaert, P. (1781) *Beschreibung zweyer merkwürdiger Fische. Neue Nordische Beyträge zur physikalischen und geographischen Erd- und Völkerbeschreibung, Naturgeschichte und Oekonomie*. St. Petersburg & Leipzig. 2, 55–57. [plates. 2, 4]
- Böhlke, E.B. (1997) Notes on the identity of elongate unpattered Indo-Pacific morays, with description of a new species (Muraenidae, Subfamily Muraeninae). *Proceedings of the Academy of Natural Sciences of Philadelphia*, 147, 89–109.
- Böhlke, E.B. & Smith, D.G. (2002) Type catalog of Indo-Pacific Muraenidae. *Proceedings of the Academy of Natural Sciences Philadelphia*, 152, 89–172.  
[http://dx.doi.org/10.1635/0097-3157\(2002\)152\[0089:TCOIPM\]2.0.CO;2](http://dx.doi.org/10.1635/0097-3157(2002)152[0089:TCOIPM]2.0.CO;2)
- Brevoort, J.C. (1856) Notes on some figures of Japanese fish taken from recent specimens by the artists of the U. S. Japan Expedition. In: Perry, M.C. (Ed.), *Narrative of the Expedition of an American Squadron to the China Seas and Japan, performed in the years 1852, 1853, and 1854 .... Vol. 2*, pp. 253–288. [plates. 3–12]
- Castle, P.H.J. (1963) The systematics, development and distribution of two eels of the genus *Gnathophis* (Congridae) in Australasian waters. *Zoological Publications from Victoria University of Wellington*, 34, 15–47.
- Castle, P.H.J. (1968) The congrid eels of the western Indian Ocean and the Red Sea. *Ichthyological Bulletin, Department of Ichthyology, Rhodes University*, 33, 685–726.
- Castle, P.H.J. & Smith, D.G. (1999) A reassessment of the eels of the genus *Bathycongrus* in the Indo-west Pacific. *Journal of Fish Biology*, 54, 973–995.  
<http://dx.doi.org/10.1111/j.1095-8649.1999.tb00851.x>
- Castle, P.H.J. & Williamson, G.R. (1975) Systematics and distribution of the eels of the *Muraenesox* group (Anguilliformes, Muraenesocidae) A preliminary report and key. *Special Publications J.L.B. Smith Institute of Ichthyology*, 15, 1–9.
- Chang, G.-C., Hung, Y.-S. & Liu, F.-K. (2011) How many fresh eel species in Taiwan? *Fishery Research Institute E-Paper*, 66, 1. [in Chinese]
- Chang, G.-C., Hung, Y.-S. & Liu, F.-K. (2013) Identification of the elver, *Anguilla celebesensis* from the surrounding waters of Taiwan. *Journal of Taiwan Fisheries Research*, 21 (1), 43–48. [in Chinese]
- Chen, C.-H. (1993) *Fishes of the Pescadores. Keelung, Taiwan*: Fisheries. Research Institute, Council of Agriculture, Executive Yuan, 397 pp. [in Chinese]
- Chen, I.-S. (2009) *Indicator species of riverine fishes in Taiwan*. Vol. 2. Diadromous fishes. National Taiwan Ocean University, Keelung, 95 pp.
- Chen, C.-H. (2004) Checklist of fishes of Penghu. *FRI special publication*, 4, 1–175.
- Chen, H.-M. & Loh, K.-H. (2007) *Gymnothorax shaoi*, a new species of moray eel (Anguilliformes, Muraenidae) from southeastern Taiwan. *Journal of Marine Science and Technology*, 15, 76–81.
- Chen, H.-M. & Shao, K.-T. (1995) New eel genus, *Cirrimaxilla*, and description of the type species, *Cirrimaxilla formosa* (Pisces, Muraenidae) from southern Taiwan. *Bulletin of Marine Science*, 57, 328–332.
- Chen, H.-M., Loh, K.-H. & Shao, K.-T. (2008) A new species of moray eel, *Gymnothorax taiwanensis*, (Anguilliformes, Muraenidae) from eastern Taiwan. *The Raffles Bulletin of Zoology*, Supplement 19, 131–134.
- Chen, H.-M., Shao, K.-T. & Chen, C.-T. (1994) A review of the muraenid eels (family Muraenidae) from Taiwan with descriptions of twelve new records. *Zoological Studies*, 33 (1), 44–64.
- Chen, H.-M., Shao, K.-T. & Chen, C.-T. (1996) A new moray eel, *Gymnothorax niphostigma*, (Anguilliformes, Muraenidae) from northern and eastern Taiwan. *Zoological Studies*, 35, 20–24.
- Chen, I.-S. & Fang, L.-S. (1999) *The freshwater and estuarine fishes of Taiwan*. National Museum of Marine Biology & Aquarium, Pingtung, 287 pp.
- Chen, I.-S., Tseng, C.-S. & Shao, K.-T. (2012) *Red data book of freshwater fishes of Taiwan*. Forestry Bureau, Council of Agriculture, Executive Yuan, Taipei, 240 pp.
- Chen, J.-P., Jan, R.-Q. & Shao, K.-T. (1997) Checklist of reef fishes from Taiping Island (Itu Aba Island), Spratly Islands, South China Sea. *Pacific Science*, 51 (2), 143–166.
- Chen, J.-P., Shao, K.-T., Jan, R.-Q., Kuo, J.-W. & Chen, J.-Y. (2010) *Marine fishes in Kenting National Park (First evised edition)*. Kenting National Park Headquarters, Pingtung, 650.
- Chen, J.T.-F. (1956) *A synopsis of the vertebrates of Taiwan*. Kaiming Press, Taipei, 292 pp. [In Chinese]
- Chen, J.T.-F. (1969) *A synopsis of the vertebrates of Taiwan. Volume 1*. Commerical Books Co., Taipei, 548 pp.
- Chen, J.T.-F. & Weng, H.T.-C. (1967) A review of the apodal fishes of Taiwan. *Biological Bulletin of Tunghai University, Ichthyology Series*, 6, 1–86.
- Chen, J.T.-F. & Yu, M.-J. (1986) *A synopsis of the vertebrates of Taiwan, revised and enlarged edition*. Commercial Press, Taipei, 1092 pp. [in Chinese]
- Chiu, Y.-C., Lin, J. & Chen, H.-M. (2013) One new record genus and three new record species of snake eels (Ophichthidae: Anguilliformes) from Taiwan. *Journal of Marine Science and Technology*, 21, supplement, 201–206.
- Chen, Y.-Y. & Mok, H.-K. (1995) *Dysomma opisthoproctus*, a new synaphobranchid eel (Pisces, Synaphobranchidae) from the northeastern coast of Taiwan. *Copeia*, 1995, 927–931.  
<http://dx.doi.org/10.2307/1447041>
- Chen, Y.-Y. & Mok, H.-K. (2001) A new synaphobranchid eel, *Dysomma longirostrum* (Anguilliformes, Synaphobranchidae), from the northeastern coast of Taiwan. *Zoological Studies*, 40, 79–83.
- Chou, T.-M. & Kao, J.-C. (2011) *The freshwater and estuarine fish of Taiwan*. Morningstar press, Taichong, 384 pp.

- Chu, K.-Y. (1957) A list of fishes from Pescadore Islands. *Report of Institute of Fishery Biology, National Taiwan University*, 1 (2), 14–23.
- Chu K.-Y. & Chen, T.-R. (1958) On the *Rhynchoconger* eels, with description of a new species. *Quarterly Journal of the Taiwan Museum*, 11, 127–129.
- Chu, Y.-T., Wu, H.-L. & Jin, X.-B. (1981) Four new species of the families Ophichthyidae and Neenchelidae. *Journal of Fisheries of China*, 5 (1), 21–27. [In Chinese, English abstract.]
- Cuvier, G. (1816) *Le Règne Animal distribué d'après son organisation pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Les reptiles, les poissons, les mollusques et les annelids*, Edition 1, v. 2, xviii + 532 pp.
- Cuvier, G. (1829) *Le Règne Animal, distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée*. Edition 2. v. 2, 406 pp.
- Ebert, D.A., Ho, H.-C., White, W.T. & de Carvalho, M.R. (2013) Introduction to the systematics and biodiversity of sharks, rays, and chimaeras (Chondrichthyes) of Taiwan. *Zooxata*, 2752 (1), 5–19.
- Eschmeyer, W.N. (2015) Catalog of fishes. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. Electronic version accessed Aug. 2015.
- Forsskål, P. (1775) *Descriptiones animalium avium, amphibiorum, piscium, insectorum, vermium; quae in itinere orientali observavit... Post mortem auctoris edidit Carsten Niebuhr*. Hauniae. 20 + xxxiv + 164 pp.
- Fourmanoir, P. & Rivaton, J. (1979) Poissons de la pente récifale externe de Nouvelle-Calédonie et des Nouvelles-Hébrides. *Cahiers de l'Indo-Pacifique*, 1 (4), 405–443.
- Fowler, H.W. (1932a) A synopsis of the fishes of China. Part III. The eels. *Hong Kong Naturalist*, 3 (1), 46–63.
- Fowler, H.W. (1932b) A synopsis of the fishes of China. Part III--concluded. The eels. *Hong Kong Naturalist*, 3 (2), 126–144.
- Fowler, H.W. (1934) Descriptions of new fishes obtained 1907 to 1910, chiefly in the Philippine Islands and adjacent seas. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 85, 233–367.
- Garman, S. (1888) On an eel from the Marshall Islands. *Bulletin of the Essex Institute*, 20, 114–116.
- Garman, S. (1899) The Fishes. In: Garman, S. (Ed.), Reports on the exploration off the west coasts of Mexico, Central and South America, and off the Galapagos islands, in charge of Alexander Agassiz by the U.S. Fish Commission Steamer "Albatross," during 1891, Lieut.-Commander Z.L. Tanner, U.S.N., commanding. XXVI. The Fishes. *Memoirs of the Museum of Comparative Zoology*, 24 (1), 1–431. [plates. 1–85 + A–M]
- Gilbert, C.H. (1891) Descriptions of apodal fishes from the tropical Pacific. In: Scientific results of explorations by the U. S. Fish Commission steamer Albatross. *Proceedings of the United States National Museum*, 14 (856), 347–352.
- Ginsburg, I. (1951) The eels of the northern Gulf Coast of the United States and some related species. *Texas Journal of Science*, 3 (3), 431–485.
- Goode, G.B. & Bean, T.H. (1879) A catalogue of the fishes of Essex County, Massachusetts, including the fauna of Massachusetts Bay and the contiguous deep waters. *Bulletin of the Essex Institute*, 11, 1–38.
- Gosline, W.A. (1951) The osteology and classification of the ophichthid eels of the Hawaiian Islands. *Pacific Science*, 5 (4), 298–320.
- Günther, A. (1870) Catalogue of the Physostomi, containing the families Gymnotidae, Symbranchidae, Muraenidae, Pegasidae, and of the Lophobranchii, Plectognathi, Dipnoi, Ganoidei, Chondropterygii, Cyclostomata, Leptocardii, in the British Museum. *Catalogue of the fishes in the British Museum*, 8, 1–549.
- Günther, A. (1872a) Report on several collections of fishes recently obtained for the British Museum. *Proceedings of the Zoological Society of London*, 1871 (pt. 3), 652–675. [plates. 53–70]
- Günther, A. (1872b) On some new species of reptiles and fishes collected by J. Brenchley, Esq. *Annals and Magazine of Natural History* (Ser. 4), 10 (60) (art. 60), 418–426.
- Günther, A. (1877) Preliminary notes on new fishes collected in Japan during the expedition of H. M. S. Challenger. *Annals and Magazine of Natural History*, (Ser. 4), 20 (119) (art. 56), 433–446.
- Günther, A. (1878) Preliminary notices of deep-sea fishes collected during the voyage of H. M. S. Challenger. *Annals and Magazine of Natural History* (Series 5), 2 (no. 7/8/9) (art. 2/22/28), 17–28, 179–187, 248–251.
- Günther, A. (1887) Report on the deep-sea fishes collected by H. M. S. Challenger during the years 1873–76. *Report on the Scientific Results of the Voyage of H. M. S. Challenger*, 22 (pt. 57), 268 pp. [66 plates]
- Hamilton, F. (1822) *An account of the fishes found in the river Ganges and its branches*. Archibald Constable & Co., Edinburgh, 405 pp.
- Hatooka, K. (1984) *Uropterygius nagoensis*, a new muraenid eel from Okinawa, Japan. *Japanese Journal of Ichthyology*, 31 (1), 20–22.
- Hatooka, K. (2002) Family Muraenidae. In: Nakabo (ed.) Fishes of Japan with pictorial keys to the species. English edition. Tokai University Press, Tokyo, pp. 196–211, pp. 1452–1455.
- Hatooka, K. & Randall, J.E. (1992) A new moray eel (*Gymnothorax*: Muraenidae) from Japan and Hawaii. *Japanese Journal of Ichthyology*, 39 (3), 183–190.
- Hatooka, K. & Yoshino, T. (1998) Two rare ophichthid species of the genus *Lamnostoma* (Pisces: Anguilliformes) from the western Pacific. *Bulletin of Osaka Museum of Natural History*, 52, 21–30.
- Herre, A. (1923) A review of the eels of the Philippine Archipelago. *Philippine Journal of Science*, 23 (2), 123–236. [plates. 1–11]

- Hibino, Y. & Kimura, S. (2015a) Revision of the *Scolecenchelys gymnota* species group with descriptions of two new species (Anguilliformes: Ophichthidae: Myrophinae). *Ichthyological Research*.  
<http://dx.doi.org/10.1007/s10228-015-0485-4>
- Hibino, Y. & Kimura, S. (2015b) A new species of *Muraenichthys* (Anguilliformes: Ophichthidae) from the Indo-Pacific, with revised generic diagnosis. *Zootaxa*, 4060 (1), 62–70.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.10>
- Hibino, Y., Ho, H.-C. & Kimura, S. (2012) A new worm eel *Neenchelys mccoskeri* (Anguilliformes: Ophichthidae) from Taiwan and Japan. *Ichthyological Research*, 59 (4), 342–346.  
<http://dx.doi.org/10.1007/s10228-012-0297-8>
- Hibino, Y., Ho, H.-C. & Kimura, S. (2015) A new genus and species of worm eels, *Sympenchelys taiwanensis* (Anguilliformes: Ophichthidae: Myrophinae) from the northwestern Pacific Ocean. *Zootaxa*, 4060 (1), 41–48.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.6>
- Ho, H.-C. & Loh, K.-H (2015) A new species of the worm-eel genus *Neenchelys* (Anguilliformes: Ophichthidae) from southern Taiwan. *Zootaxa*, 4060 (1), 52–55.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.8>
- Ho, H.-C. & Shao, K.-T. (2010) A remarkable new record of congrid eel *Blachea xenobranchialis* Karrer and Smith, 1980 (Pisces: Anguilliformes) from Taiwan. *Journal of Fishery Society of Taiwan*, 37 (4), 285–290.
- Ho, H.-C. & Shao, K.-T. (2011) Annotated checklist and type catalog of fish genera and species described from Taiwan. *Zootaxa*, 2957, 1–74.
- Ho, H.-C., McCosker, J.E. & Smith D.G. (2013) Revision of the worm eel genus *Neenchelys* (Ophichthidae; Myrophinae), with descriptions of three new species from the western Pacific Ocean. *Zoological Studies*, 52, 1–20.  
<http://dx.doi.org/10.1186/1810-522X-52-58>
- Ho, H.-C., McCosker, J.E. & Smith D.G. (2015c) Renaming of three recently described eels of the genus *Neenchelys* (Teleostei: Anguilliformes: Ophichthidae) from the western Pacific. *Zootaxa*, 4060 (1), 49–51.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.7>
- Ho, H.-C., Smith D.G. & Shao, K.-T. (2015b) Notes on the congrid eel genus *Parabathymyrus* from the western Pacific Ocean, with the description of a new species (Pisces: Anguilliformes: Congridae). *Zootaxa*, 4060 (1), 131–139.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.15>
- Ho, H.-C., Tighe, K. & Smith, D.G. (2015d) Review of the arrowtooth eel genera *Dysomma* and *Dysommmina* in Taiwan, with the description of a new species (Anguilliformes: Synaphobranchidae: Ilyphinae). *Zootaxa*, 4060 (1), 86–104.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.12>
- Ho, H.-C., McCosker, J.E., Smith D.G. & Shao, K.-T. (2015a) Introduction to the systematics and biodiversity of eels (orders Anguilliformes and Saccopharyngiformes) of Taiwan. *Zootaxa*, 4060 (1), 5–18.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.3>
- Ho, H.-C., Smith, D.G., Wang, S.-I., Shao, K.-T., Ju, Y.-M. & Chang, C.-W. (2010) Specimen catalog of Pisces collection of National Museum of Marine Biology and Aquarium transferred from Tinghai University and Aquarium. (II) Order Anguilliformes. *Platax*, 7, 13–34.
- International Commission on Zoological Nomenclature (2012) Amendment of Articles 8, 9, 10, 21 and 78 of the International Code of Zoological Nomenclature to expand and refine methods of publication. *Zootaxa*, 3450, 1–7.
- Jenkins, O.P. (1903) Report on collections of fishes made in the Hawaiian Islands, with descriptions of new species. *Bulletin of the U. S. Fish Commission*, 22, 415–511.
- Johnson, J.Y. (1862) Descriptions of some new genera and species of fishes obtained at Madeira. *Proceedings of the Zoological Society of London*, 1862, 167–180.
- Jordan, D.S. & Evermann, B.W. (1902) Notes on a collection of fishes from the island of Formosa. *Proceedings of the United States National Museum*, 25, 315–368.  
<http://dx.doi.org/10.5479/si.00963801.25-1289.315>
- Jordan, D.S. & Richardson, R.E. (1908) Fishes from islands of the Philippine Archipelago. *Bulletin of the Bureau of Fisheries*, 27, 233–287.
- Jordan, D.S. & Richardson, R.E. (1909) A catalogue of the fishes of the island of Formosa, or Taiwan, based on the collections of Dr. Hans Sauter. *Memoirs of the Carnegie Museum*, 4, 159–204.
- Jordan, D.S. & Seale, A. (1906) The fishes of Samoa. Description of the species found in the archipelago, with a provisional check-list of the fishes of Oceania. *Bulletin of the Bureau of Fisheries*, 25, 173–455.
- Jordan, D.S. & Snyder, J.O. (1900) A list of fishes collected in Japan by Keinosuke Otaki, and by the United States steamer Albatross, with descriptions of fourteen new species. *Proceedings of the United States National Museum*, 23 (1213), 335–380.  
<http://dx.doi.org/10.5479/si.00963801.23-1213.335>
- Jordan, D.S. & Snyder, J.O. (1901) A review of the apodal fishes or eels of Japan, with descriptions of nineteen new species. *Proceedings of the United States National Museum*, 23 (1239), 837–890.  
<http://dx.doi.org/10.5479/si.00963801.23-1239.837>
- Jordan, D.S. & Snyder, J.O. (1904) Notes on collections of fishes from Oahu Island and Laysan Island, Hawaii, with descriptions of four new species. *Proceedings of the United States National Museum*, 27 (1377), 939–948.  
<http://dx.doi.org/10.5479/si.00963801.27-1377.939>

- Jordan, D.S., Tanaka, S. & Snyder, J.O. (1913) A catalogue of the fishes of Japan. *Journal of the College of Science, Imperial University, Tokyo*, 33 (1), 1–497.
- Kamohara, T. (1938) *On the offshore bottom-fishes of Prov. Tosa, Shikoku, Japan*. Maruzen Kobushiki Kaisha, Tokyo, 86 pp.
- Kanazawa, R.H. (1958) A revision of the eels of the genus *Conger* with descriptions of four new species. *Proceedings of the United States National Museum*, 108 (3400), 219–267.
- Karmovskaya, E.S. (1985) Mesopelagic eels of family Derichthyidae. *Journal of Ichthyology*, 25 (6), 119–134.
- Karmovskaya, E.S. (1994) Systematics and distribution of the eel genus *Gavialiceps* (Congridae) in the Indo-west Pacific. *Journal of Ichthyology*, 34 (3), 73–89.
- Karmovskaya, E.S. (2004) Benthopelagic bathyal conger eels of families Congridae and Nettastomatidae from the western tropical Pacific, with descriptions of ten new species. *Journal of Ichthyology*, 44 (Supplement 1), S1–32.
- Karrer, C. (1983) Anguilliformes du Canal de Mozambique (Pisces, Teleostei). *Faune Tropicale*, 23, 1–116.
- Karrer, C. & Smith, D.G. (1980) A new genus and species of congrid eel from the Indo-west Pacific. *Copeia*, 1980 (4), 642–648.  
<http://dx.doi.org/10.2307/1444440>
- Kaup, J.J. (1856a) Uebersicht der Aale. *Archiv für Naturgeschichte*, 22 (1), 41–77.
- Kaup, J.J. (1856b) *Catalogue of the apodal fish, in the collection of the British Museum*. British Museum, London, viii + 163 pp.
- Kaup, J.J. (1859) Neue aalaehnliche Fische des Hamburger Museums, beschrieben und abgebildet. *Abhandlungen aus dem Gebiete der Naturwissenschaften herausgegeben von dem Naturwissenschaftlichen Verein in Hamburg*, 4 (2), 1–29 + supplement. 1–4.
- Klausewitz, W. & Eibl-Eibesfeldt, I. (1959) Neue Röhrenaale von den Maldiven und Nikobaren (Pisces, Apodes, Heterocongridae). *Senckenbergiana Biologica*, 40 (3/4), 135–153.
- Klunzinger, C.B. (1871) Synopsis der Fische des Rothen Meeres. II. Theil. Verhandlungen der K.-K. zoologisch-botanischen Gesellschaft in Wien, 21, 441–688.
- Kuroda, N. (1947) Additions to the catalogue of the fishes of Shizuura, Suruga and the vicinity, (VIII). *Seibutu, Supplemental* 1, 25–31.
- Lacepède, B.G.E. (1803) *Histoire naturelle des poissons*. v. 5: lxviii + 803 pp.
- Lay, G.T. & Bennett, E.T. (1839) Fishes. In: Beechey, F.W. (ed.) *The zoology of Captain Beechey's voyage; compiled from the collections and notes made by Captain Beechey, the officers and naturalist of the expedition, during a voyage to the Pacific and Behring's Straits performed in His Majesty's ship Blossom, under the command of Captain F.W. Beechey, R. N., F.R.S., &c. &c., in the years 1825, 26, 27 and 28*, H. G. Bohn, London, pp. 41–75.
- Lee, S.-C. (1967) Report on the fishes of Pescadore Islands. *The Biological Bulletin of Taiwan Normal University*, 29 (1), 53–55.
- Lee S.-C. (1980) New records of fishes from Lanyu (Botel Tobago), Taiwan. *Bulletin of the Institute of Zoology, Academia Sinica*, 19 (2), 57–61.
- Lee, S.-C. & Tseng, W.-N. (1985) First record of the moray eel, *Gymnothorax chlamydatus* from Taiwan. *Journal of Taiwan Museum*, 24 (2), 295–296.
- Lee, S.-C. & Yang, H.-C. (1966) Notes on the congrid eels of Taiwan. *Bioformosa*, 1, 52–63.
- Lesson, R.P. (1828) Description du nouveau genre *Ichthyophis* et de plusieurs espèces inédites ou peu connues de poissons, recueillis dans le voyage autour du monde de la Corvette "La Coquille". *Mémoires de la Société d'Histoire Naturelle Paris*, 4, 397–412.
- Li, L.-K. (1960) Additions to the apodal ichthyofauna of Taiwan (Formosa). *Journal of Taiwan Museum*, 13 (1–2), 83–89.
- Liang, Y.-S. & Yeh, C.-F. (1964) List of the fish specimens preserving in the Museum of the Department of Zoology National Taiwan University. *Occasional Paper, Department of Zoology, National Taiwan University*, 1–67.
- Liaw, W.-K. (1960) Study on the fishes from Ma-kung, Pescadores Island (Penghu). *Report of Laboratory of Fishery Biology, Taiwan Fisheries Research Institute*, 13, 1–26.
- Lin, C.-J. (2007) *Freshwater fishes and shrimps of Taiwan*. Commonwealth Publishing Group, Taipei, 239 pp. [in Chinese]
- Lin, J., Shao, K.-T. & Chen, H.-M. (2013) Taxonomic study of pike congers (Anguilliformes: Muraenesocidae), with identification of muraenesocid collections in Taiwan. *Journal of Marine Science and Technology*, 21 (Supplement), 166–174.
- Lin, J., Smith, D.G., Shao, K.-T. & Chen, H.-M. (2015) *Saurenchelys gigas* sp. nov., a new nettastomatid eel (Teleostei, Anguilliformes, Nettastomatidae) from the western central Pacific. *Zootaxa*, 4060 (1), 121–130.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.14>
- Loh, K.-H., Shao, K.-T. & Chen, H.-M. (2011) *Gymnothorax melanosomatus*, a new morey eel (Teleostei: Anguilliformes: Muraenidae) from southeastern Taiwan. *Zootaxa*, 3134, 43–52.
- Loh, K.-H., Chen, I.-S., Randall, J. E. & Chen H.-M. (2008) A review and molecular phylogeny of the moray eel subfamily Uropterygiinae (Anguilliformes: Muraenidae) from Taiwan, with description of a new species. *Raffles Bulletin of Zoology, Supplement 19*, 135–150.
- Loh, K.-H., Shao, K.-T. & Chen, H.-M. (2012) Additions to the Taiwan marine eel fauna with first records of three rare moray eels (Anguilliformes: Muraenidae). *Journal of Marine Science and Technology*, 20 (2), 210–215.
- Loh, K.-H., Shao, K.-T., Chong, V.-C. & Chen, H.-M. In: press. Additions to the Taiwan eel fauna with five newly recorded

- species of moray eels (Anguilliformes: Muraenidae), and redescription of *Gymnothorax sagmacephalus*. *Journal of Marine Science and Technology*.
- Loh, K.-H., Shao, K.-T., Ho, H.-C., Lim, P.-E. & Chen, H.-M. (2015) A new species of moray eel (Anguilliformes: Muraenidae) from Taiwan, with comments on related elongate unpatterned species. *Zootaxa*, 4060 (1), 30–40.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.5>
- Machida, Y. & Ohta, S. (1993) New record for *Neenchelys daedalus* (Ophichthidae) from Japan. *Japanese Journal of Ichthyology*, 39 (4), 391–394.
- Matsubara, K. (1955) *Fish morphology and hierarchy. Part 1*. Ishizaki-Shoten, Tokyo, 789 pp. [In Japanese]
- Matsubara, K. & Asano, H. (1959) A new eel of the genus *Kaupichthys*. *Copeia*, 1959 (4), 293–297.  
<http://dx.doi.org/10.2307/1439885>
- Matsubara, K. & Ochiai, A. (1951) On the conger eels related to *Arisoma nystromi* (Jordan et Snyder) found in the waters of Japan and China. *Memoirs of the College of Agriculture Kyoto University*, 59, 1–18.
- McClelland, J. (1843) Description of a collection of fishes made at Chusan and Ningpo in China, by Dr. G. R. Playfair, Surgeon of the Phlegethon, war steamer, during the late military operations in that country. *Calcutta Journal of Natural History*, 4 (4), 390–41. [plates. 21–25]
- McClelland, J. (1844) Apodal fishes of Bengal. *Calcutta Journal of Natural History*, 5 (18), 151–226. [plates. 5–14]
- McCosker, J.E. (1982) A new genus and two new species of remarkable Pacific worm eels (Ophichthidae, subfamily Myrophinae). *Proceedings of the California Academy of Sciences*, 43, 59–66.
- McCosker, J.E. (2014) A gigantic deepwater worm eel (Anguilliformes: Ophichthidae) from the Verde Island Passage, Philippine Archipelago. In: Williams, G.C. & Gosliner, T.M. (Eds.), *The Coral Triangle. The 2011 Hearst Philippine Biodiversity Expedition*. California Academy of Sciences, San Francisco, pp 333–340.
- McCosker, J.E. & Chen, Y.-Y. (2000) A new species of deepwater snake-eel, *Ophichthus aphotistos*, with comments on *Neenchelys retropinna* (Anguilliformes: Ophichthidae) from Taiwan. *Ichthyological Research*, 47 (4), 353–357.  
<http://dx.doi.org/10.1007/BF02674262>
- McCosker, J.E. & Ho, H.-C. (2014) Redescription of the sharp-fanged snake eel *Brachysomophis longipinnis* (Anguilliformes: Ophichthidae). *Platax*, 11, 95–101.
- McCosker, J. E. & Ho, H.-C. (2015) New species of the snake eels *Echelus* and *Ophichthus* (Anguilliformes: Ophichthidae) from Taiwan. *Zootaxa*, 4060 (1), 71–85.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.11>
- McCosker, J.E., Chen, W.-L. & Chen, H.-M. (2009) Comments on the snake-eel genus *Xyrias* (Anguilliformes: Ophichthidae) with the description of a new species. *Zootaxa*, 2289, 61–67.
- McCosker, J. E., Loh, K.-H., Lin, J. & Chen, H.-M. (2012) *Pylorobranchus hoi*, a new genus and species of myrophine worm-eel from Taiwan (Anguilliformes: Ophichthidae). *Zoological Studies*, 51 (7), 1188–1194.
- McCosker, J.E. & Chen, Y.-Y. (2000) A new species of deepwater snake-eel, *Ophichthus aphotistos*, with comments on *Neenchelys retropinna* (Anguilliformes, Ophichthidae) from Taiwan. *Ichthyological Research*, 47, 353–357.  
<http://dx.doi.org/10.1007/BF02674262>
- McCosker, J.E., Ide, S. & Endo, H. (2012) Three new species of ophichthid eels (Anguilliformes: Ophichthidae) from Japan. *Bulletin of the National Museum of Nature and Science (Ser. A)* Supplement 6, 1–16.
- McCosker, J.E. & Randall, J.E. (1993) Finless snake-eels of the genus *Cirricaecula* (Anguilliformes, Ophichthidae), with the description of *C. macdowelli* from Taiwan. *Japanese Journal of Ichthyology*, 40, 189–192.
- McCosker, J.E. & Randall, J.E. (2001) Revision of the snake-eel genus *Brachysomophis* (Anguilliformes, Ophichthidae), with description of two new species and comments on the species of *Mystriophis*. *Indo-Pacific Fishes*, 33, 1–32.
- McCosker, J.E. & Randall, J.E. (2002) *Ophichthys melanochir* a junior synonym of the Highfin Snake Eel *Ophichthys altipennis*. *Copeia*, 2002 (3), 798–799.
- McCosker, J.E. & Rosenblatt, R.H. (1993) A revision of the snake eel genus *Myrichthys* (Anguilliformes: Ophichthidae) with the description of a new eastern Pacific species. *Proceedings of the California Academy of Sciences*, 48, 153–169.
- McCosker, J.E., Hatooka, K., Ohnishi, N. & Endo, H. (2011) Redescription and designation of a neotype for *Aphthalmichthys kuro* Kuroda 1947, and its placement in *Callechelys* (Anguilliformes: Ophichthidae). *Ichthyological Research* v. 58, 272–277.  
<http://dx.doi.org/10.1007/s10228-011-0233-3>
- Mok, H.-K., Lee, C.-Y. & Chan, H.-J. (1991) *Meadia roseni*, a new synaphobranchid eel from the coast of Taiwan (Anguilloidea, Synaphobranchidae). *Bulletin of Marine Science*, 48, 39–45.
- Nielsen, J.G. & Smith, D.G. (1978) The eel family Nemichthyidae (Pisces: Anguilliformes). *Dana Report*, 88, 1–71. [plates. 1–2]
- Norman, J.R. (1922) Two new fishes from New Britain and Japan. *Annals and Magazine of Natural History* (Series 9), 10 (56), 217–218.  
<http://dx.doi.org/10.1080/00222932308632764>
- Okamura, O. & Kitajima, T. (Eds.) (1984) *Fishes of the Okinawa Trough and the adjacent waters. Vol. 1. The intensive research of unexploited fishery resources on continental slopes*. Japan Fisheries Resource Conservation Association, Tokyo. v. 1, 1–414. [plates. 1–205]
- Ogilby, J.D. (1895) Description of a new Australian eel. *Proceedings of the Linnean Society of New South Wales* (Series 2), 9

- (4) (for 1894), 720–721.
- Ogilby, J.D. (1897) Some new genera and species of fishes. *Proceedings of the Linnean Society of New South Wales*, 22 (2), 245–251.
- Okada, Y. (1938) *A catalogue of vertebrates of Japan*. Maruzen, Tokyo, 412 pp.
- Okada, Y. & Matsubara, K. (1938) *Keys to the fishes and fish-like animals of Japan*. Sanseido, Tokyo, 584 pp. [in Japanese]
- Oshima, M. (1919) Contributions to the study of the fresh water fishes of the island of Formosa. *Annals of the Carnegie Museum*, 12 (2–4), 169–328.
- Playfair, R.L. & Günther, A. (1867) *The fishes of Zanzibar, with a list of the fishes of the whole east coast of Africa*. John Van Voorst, London, xiv + 153 pp.
- Quoy, J.R.C. & Gaimard, J.P. (1824) Description des Poissons. Chapter IX. In: Freycinet, L. de, *Voyage autour du Monde...exécuté sur les corvettes de L. M. "L'Uranie" et "La Physicienne," pendant les années 1817, 1818, 1819 et 1820*. Paris, pp. 192–401.
- Randall, J.E., Allen, G.R. & Steene, R.C. (1997) *Fishes of the Great Barrier Reef and Coral Sea*. Revised edition. Crawford House Press, Bathurst NSW, 557 pp.
- Regan, C.T. (1905) On a collection of fishes from the inland sea of Japan made by Mr. R. Gordon Smith. *Annals and Magazine of Natural History* (Series 7), 15 (85) (art. 2), 17–26.
- Regan, C.T. (1909) Descriptions of new marine fishes from Australia and the Pacific. *Annals and Magazine of Natural History* (Series 8), 4 (23), 438–440.  
<http://dx.doi.org/10.1080/00222930908692695>
- Richardson, J. (1845a) Ichthyology. Part 2. In Hinds, R.B. (Ed.), *The zoology of the voyage of H. M. S. Sulphur, under the command of Captain Sir Edward Belcher, R. N., C. B., F. R. G. S., etc., during the years 1836–42*, No. 9. London: Smith, Elder & Co., pp. 71–98. [plates. 45–54]
- Richardson, J. (1845b) Ichthyology. Part 3. In: Hinds, R.B. (Ed.), *The zoology of the voyage of H. M. S. Sulphur, under the command of Captain Sir Edward Belcher, R. N., C. B., F. R. G. S., etc., during the years 1836–42*, No. 10. London: Smith, Elder & Co., pp. 99–150. [plates. 55–64]
- Richardson, J. (1848a) Fishes. In: A. Adams (Ed.). *The zoology of the voyage of H. M. S. Samarang; under the command of Captain Sir Edward Belcher, during the years 1843–1846*. Reeve & Benham, London, 128 pp. [10 plates]
- Richardson, J. (1848b) *Ichthyology of the voyage of H.M.S. Erebus & Terror, under the command of Captain Sir James Clark Ross, R.N., F.R.S. Edward Newman*, London, viii + 139 pp.
- Robins, C.H. & Robins, C.R. (1976) New genera and species of dysommine and synaphobranchine eels (Synaphobranchidae) with an analysis of the Dysommidae. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 127 (18), 249–280.
- Robins, C.H. & Robins, C.R. (1989) Family Synaphobranchidae. In: Böhlke, E.B. (Ed.), Volume One: Orders Anguilliformes and Saccopharyngiformes. Fishes of the western North Atlantic. *Memoirs of the Sears Foundation of Marine Research*, 9, 207–253.
- Rüppell, W.P.E.S. (1828–30) *Atlas zu der Reise im nördlichen Africa. Fische des Rothen Meeres*. Frankfurt-am-Main, 141 + 3 pp. [35 col. plates.]
- Sabaj Pérez, M.E. (2014) Standard symbolic codes for institutional resource collections in herpetology and ichthyology: an Online Reference. Version 5.0 (22 September 2014). Available from: <http://www.asih.org/>, American Society of Ichthyologists and Herpetologists, Washington, DC.
- Saldanha, L. & Quero, J.-C. (1994) *Channomuraena bauchotae* (Anguilliformes: Muraenidae), nouvelle espèce de l'Île de la Réunion, Océan Indien. *Cybium*, 18 (3), 307–313.
- Sasaki, K. & Amaoka, K. (1991) *Gymnothorax prolatus*, a new moray from Taiwan. *Japanese Journal of Ichthyology*, 38 (1), 7–10.
- Schmidt, J. (1928) The fresh-water eels of Australia with some remarks on the short-finned species of Anguilla. *Records of the Australian Museum*, 16 (4), 179–210.  
<http://dx.doi.org/10.3853/j.0067-1975.16.1928.785>
- Schmidt, J. (1931) Oceanographic expedition of the Dana, 1928–1930. *Nature (London)*, 127, 444–446, 487–490.  
<http://dx.doi.org/10.1038/127487a0>
- Schultz, L.P. (1943) Fishes of the Phoenix and Samoan islands collected in 1939 during the expedition of the U. S. S. "Bushnell." *Bulletin of the United States National Museum*, 180, x + 316 pp.
- Schultz, L.P. & Woods, L.P. (1949) Keys to the genera of echelid eels and the species of Muraenichthys of the Pacific, with two new species. *Journal of the Washington Academy of Sciences*, 39 (5), 169–174.
- Schultz, L.P., Herald, E.S., Lachner, E.A., Welander, A.D. & Woods, L.P. (1953) *Fishes of the Marshall and Marianas islands*. Vol. I. Families from Asymmetrontidae through Siganidae, *Bulletin of the United States National Museum*, 202, xxxii + 685 pp.
- Seale, A. (1906) Fishes of the South Pacific. *Occasional Papers of the Bernice Pauahi Bishop Museum of Polynesian Ethnology and Natural History*, 4 (1), 1–89.
- Seale, A. (1917) New species of apodal fishes. *Bulletin of the Museum of Comparative Zoology*, 61 (4), 79–94.
- Séret, B., Brischoux, F., Bonnet, X. & Shine R. (2008) First record of *Cirrimaxilla formosa* (Muraenidae) from New Caledonia, found in sea snake stomach contents. *Cybium*, 32 (2), 191–192.
- Shao, K.-T. (1990) Garden eels from Taiwan, with description of a new species. *Uo (Japanese Society of Ichthyologists)*, 40, 1–

- Shao, K.-T. (2015) Taiwan Fish Database. <http://fishdb.sinica.edu.tw> (Jul. 2015).
- Shao K.-T., Chen, J.-P., Ho, L.-T., Lin, C.-P., Kao, P.-H., Lin, P.-L. & Chen, L.-S. (1994a) Checklist and distributional pattern of the fishes of the Pescadores Islands. In: Faculty of Fisheries, Kasetsart University (Ed.) *Proceeding of the 4th Indo-Pacific fish Conference: systematics and evolution of Indo-Pacific fishes*. Kasetsart University, Bangkok, pp. 267–280.
- Shao, K.-T., Lin, P.-L., Chen, L.-S. & Hsieh, L.-Y. (1994b) Catalogue of the fish specimens held in the National Marine Science Museum (1). *Acta Zoologica Taiwanica*, 5 (1), 47–76.
- Shao, K.-T., Ho, H.-C., Lin, P.-L., Lee, P.-F., Lee, M.-Y., Tsai, C.-Y., Liao, Y.-C. & Lin, Y.-C. (2008) A checklist of the fishes of southern Taiwan, Northern South China Sea. *Raffles Bulletin of Zoology*, Supplement 19, 233–271.
- Shaw, G. & Nodder, F.P. (1795) The Naturalist's Miscellany, or coloured figures of natural objects; drawn and described from nature. London, unnum. pages.
- Shaw, G. & Nodder, F.P. (1797) The Naturalist's Miscellany, or coloured figures of natural objects; drawn and described from nature. London, unnum. pages.
- Shen S.-C. (1974) Two new records of the Genus *Rhinomuraena* (Family Muraenidae) found in the waters of Taiwan. *Acta Oceanographica Taiwanica*, 4, 181–190.
- Shen, S.-C. (1984a) *Synopsis of fishes of Taiwan*. Southern Materials Center, Taipei, Taiwan, 533 pp.
- Shen, S.-C. (1984b) *Coastal fishes of Taiwan*. National Taiwan Museum, Taipei, Taiwan, 189 pp.
- Shen, S.-C. (1988) Catalogue of the fish specimen deposited in the museum of the Department of Zoology, National Taiwan University. *Acta Zoologica Taiwanica*, 2, 1–162.
- Shen, S.-C. (1998) A review of congrid eels of the genus *Ariosoma* from Taiwan, with description of a new species. *Zoological Studies*, 37 (1), 7–12.
- Shen, S.-C. & Wu, K.-Y. (2011) *Fishes of Taiwan*. National Museum of Marine Biology & Aquarium, Pingtung, 895 pp.
- Shen, S.-C., Lee, S.-C., Shao, K.-T., Mok, H.-K., Chen, C.-T. & Chen, C.-H. (1993) *Fishes of Taiwan*. Department of Zoology, National Taiwan University, Taipei, 960 pp.
- Smith, D.G. (1999a) Synaphobranchidae, Cutthroat eels. In: Carpenter, K.E. & Niem, V.H. (Eds.), *Species identification guide for fisheries purposes. The living marine resources of the western central Pacific. Volume 3. Batoid fishes, chimeras and bony fishes part 1 (Elopidae to Linophrynidae)*. FAO, Rome, pp. 1658–1661.
- Smith, D.G. (1999b) Congridae, Conger eels. In: Carpenter, K.E. & Niem, V.H. (Eds.), *Species identification guide for fisheries purposes. The living marine resources of the western central Pacific. Volume 3. Batoid fishes, chimeras and bony fishes part 1 (Elopidae to Linophrynidae)*. FAO, Rome, pp. 1680–1687.
- Smith, D.G. (2000) Family Congridae. In: Randall, J.E. & Lim, K.K.P. (Eds.), A checklist of the fishes of the South China Sea. *Raffles Bulletin of Zoology* Supplement 8, 1–586.
- Smith, D.G. (2012) A checklist of the moray eels of the world (Teleostei: Anguilliformes: Muraenidae). *Zootaxa*, 3474, 1–64.
- Smith, D.G. & Böhlke, J.E. (1983) *Neenchelys retropinna*: a new worm eel (Pisces: Ophichthidae) from the Indian Ocean. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 135, 80–84.
- Smith, D.G., Böhlke, J.E. & Castle, P.H.J. (1981) A revision of the nettastomatid eel genera *Nettastoma* and *Nettenchelys* (Pisces: Anguilliformes), with descriptions of six new species. *Proceedings of the Biological Society of Washington*, 94 (2), 535–560.
- Smith, D.G. & Kanazawa, R.H. (1977) Eight new species and a new genus of congrid eels from the western north Atlantic with redescriptions of *Ariosoma analis*, *Hildebrandia guppyi*, and *Rhechias vicinalis*. *Bulletin of Marine Science*, 27 (3), 530–543.
- Smith, D.G., Lin, J., Chen, H.-M. & Pogonoski, J.J. (2015) The eels of the genus *Nettenchelys*, with description of a new species from Taiwan (Teleostei: Anguilliformes: Nettastomatidae). *Zootaxa*, 4060 (1), 105–120. <http://dx.doi.org/10.11646/zootaxa.4060.1.13>
- Smith, J.L.B. (1962) The moray eels of the western Indian Ocean and the Red Sea. *Ichthyological Bulletin, Department of Ichthyology, Rhodes University*, 23, 421–444. [plates. 53–62]
- Smith, J.L.B. (1965) The Indian genus *Bathymyrus* Alcock, 1889 with description of a new species from Vietnam. *Occasional Papers of the Department of Ichthyology, Rhodes University*, 2, 1–11. [1 plate]
- Snyder, J.O. (1904) A catalogue of the shore fishes collected by the steamer "Albatross" about the Hawaiian Islands in 1902. *Bulletin of the U. S. Fish Commission*, 22, 513–538. [plates. 1–13]
- Snyder, J.O. (1908) Descriptions of eighteen new species and two new genera of fishes from Japan and the Riu Kiu Islands. *Proceedings of the United States National Museum*, 35 (1635), 93–111.
- Tan, H.J. (1983) New records of fishes from China Sea. *Investigatio et studium naturae*, 1983, 25–26. [in Chinese]
- Tanaka, S. (1911) *Figures and descriptions of the fishes of Japan, including Riukiu Islands, Bonin Islands, Formosa, Kurile Islands, Korea and southern Sakhalin*, 2, 19–34. [plates. 6–10]
- Tanaka, S. (1917) *Figures and descriptions of the fishes of Japan including Riukiu Islands, Bonin Islands, Formosa, Kurile Islands, Korea and southern Sakhalin*. 26, 455–474. [plates. 126–130]
- Tanaka, S. (1918) Two new species of Japanese fishes. *Dobutsugaku Zasshi = Zoological Magazine Tokyo*, 30 (352), 51–52.
- Temminck, C.J. & Schlegel, H. (1846) Pisces. In: *Fauna Japonica, sive descriptio animalium quae in itinere per Japoniam suscepto annis 1823–30 collegit, notis observationibus et adumbrationibus illustravit P. F. de Siebold*. Part 1 (10–14), 173–269.

- Tighe, K., Pogonoski, J.J., Hibino, Y., Ho, H.-C. & Nguyen, Q.V. (2015) Three new species of the genus *Chlopsis* (Anguilliformes: Chlopsidae) from the Indo-Pacific. *Zootaxa*, 4060 (1), 19–29.  
<http://dx.doi.org/10.11646/zootaxa.4060.1.4>
- Tseng, W.-N. (1982) On the new record of *Anguilla celebesensis* from Taiwan. *Chinese Bioscience*, 19, 57–66. [in Chinese]
- Tseng, W.-N. & Tabeta, O. (1983) First record of the short-finned eel *Anguilla bicolor pacific* elvers from Taiwan. *Bulletin of the Japanese Society for the Science of Fish*, 49, 27–32.  
<http://dx.doi.org/10.2331/suisan.49.27>
- Vaillant, L.L. (1882) Sur un poisson des grandes profondeurs de l'Atlantique, l'*Eurypharynx pelecanoides*. *Comptes Rendus Hebdomadaires des séances de l'Académie des sciences*, 95, 1226–1228.
- Vaillant, L.L. (1888) *Expéditions scientifiques du "Travailleur" et du "Talisman" pendant les années 1880, 1881, 1882, 1883. Poissons*. Masson, Paris, 406 pp.
- Watanabe, S., Aoyama, J. & Tsukamoto, K. (2009) A new species of freshwater eel *Anguilla luzonensis* (Teleostei: Anguillidae) from Luzon Island of the Philippines. *Fisheries Science (Tokyo)*, 75, 387–392.  
<http://dx.doi.org/10.1007/s12562-009-0087-z>
- Whitley, G.P. (1927) The fishes of Michaelmas Cay, North Queensland. *Records of the Australian Museum*, 16 (1), 1–32.  
<http://dx.doi.org/10.3853/j.0067-1975.16.1927.776>