



ZHI-TENG CHEN (2020) First discovery of *Leuctra fusca tergostyla* Wu, 1973 in Sichuan Province of China and notes on its taxonomic status, with new illustrations for *Rhopalopssole sinensis* Yang & Yang, 1993 (Plecoptera: Leuctridae). *Zootaxa*, 4929: 195–206.

- 1) The female of *Rhopalopssole sinensis* was not illustrated for the first time in this contribution. Fig. 23.12 in Sivec *et al.* (2008) may refer to this species, and later the female was described by Li *et al.* (2017). These authors indicated that the distinctive character of this species is the long apical whorl of hairs on antennae.
- 2) The long apical whorl of hairs on antennae was also a distinct character in Fujian specimens of *R. sinensis*, see Fig. 12A.
- 3) In Fig. 1, the distribution of *Leuctra fusca* (Linnaeus, 1758) should also cover Lebanon, but not Crete, Cyprus and only the Elburz Range (Caspian shore) of Iran.
- 4) In remarks regarding *L. fusca tergostyla*, a citation for the morphology of the nominal subspecies should include Lillehammer (1988).
- 5) Qian *et al.* (2014) was incorrectly cited as Qian & Du (2014) in the text. In references, add Stephens, J.F. (1836) Family II.—Perlidae, Leach. *Illustrations of British Entomology*, 6, 134–145.

Additional References

- Li, W.H., Murányi, D., Gamboa, M., Yang, D. & Watanabe, K. (2017) New species and records of Leuctridae (Plecoptera) from Guangxi, China, on the basis of morphological data, with emphasis on *Rhopalopssole*. *Zootaxa*, 4243 (1), 165–176.
<https://doi.org/10.11646/zootaxa.4243.1.8>
- Lillehammer, A. (1988) Stoneflies (Plecoptera) of Fennoscandia and Denmark. *Fauna Entomologica Scandinavica*, 21, 1–165